



HAL
open science

Methylation analysis in monocytes at postpartum period in dairy cattle

Hélène Jammes, Maxime Gasselín, Jean-Philippe Perrier, François Piumi, Luc
Jouneau, Hala Al Hadami, Audrey Prézélin, Marion Boutinaud, Christine
Leroux, José Pires, et al.

► **To cite this version:**

Hélène Jammes, Maxime Gasselín, Jean-Philippe Perrier, François Piumi, Luc Jouneau, et al.. Methylation analysis in monocytes at postpartum period in dairy cattle. PAG XXV - Plant and Animal Genome Conference, Jan 2017, San Diego, United States. , 2017. hal-02737532

HAL Id: hal-02737532

<https://hal.inrae.fr/hal-02737532v1>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



The Largest Ag-Genomics Meeting in the World.

PLANT & ANIMAL GENOME CONFERENCE XXIV

www.intlpag.org

January 14-18, 2017 San Diego, CA

P1059: Methylation Analysis in Monocytes at Postpartum Period in Dairy Cattle

Epigenetic modifications such as DNA methylation play a role in regulating gene expression and consequently in biological processes, such as those involved in health and disease. In dairy cows, profound changes occur during the post calving period, leading to immunosuppression and increased susceptibility to diseases. In order to monitor the postpartum health status of cows, one challenge is to describe the methylome of a purified subpopulation of immune cells, such as monocytes, and to determine its alterations/modifications in response to environmental or physiological changes. In this study, genome-wide DNA methylation profiles were obtained from purified monocytes (n=11), Peripheral Blood Mononuclear Cells (PBMC; n=4) sampled at D15 after calving and fibroblasts (n=2) using Reduced Representation Bisulfite Sequencing (RRBS). After sequencing and mapping to the reference genome (UMD3.1 assembly), 34.6% to 40.2% of uniquely mapped reads were obtained. Only CpGs covered between 10X and 500X (CpG₁₀₋₅₀₀), were analyzed, which represented 55 ± 9.8 % of total CpGs with no significant differences between libraries. The chromosomal distribution of CpG₁₀₋₅₀₀ was independent of the chromosome length (pvalue=0.48) but significantly associated with the coding gene content (pvalue=0.0001076). The global CpG₁₀₋₅₀₀ methylation scores varied between cell types (fibroblasts: $47\% \pm 0.1$, monocytes: $54.5\% \pm 2.9$ and PBMC: $54.1\% \pm 1.8$), and 19417 monocyte-specific Differentially Methylated Cytosines (DMCs) corresponding to 1572 DMRs were identified. This study therefore highlights CpGs and regions displaying a specific methylation pattern in monocytes, which could be targeted by epigenetic changes induced by environmental conditions (husbandry, nutrition, infection challenges...).

Authors

Helene Jammes

UMR BDR, INRA, ENVA,
Université Paris Saclay

Marion Boutinaud

INRA

Maxime Gasselin

INRA

Christine Leroux

INRA

Jean Philippe Perrier

UMR BDR, INRA, ENVA,
Université Paris Saclay

José Pires

INRA

François Piumi

INRA

Michael Weber

CNRS

Luc Jouneau

UMR BDR, INRA, ENVA,
Université Paris Saclay

Bruno Pount

Auriva-Elevage

Hala Al Adhami

CNRS

Jackie Jzawadzki

Pilardière group

Audrey Prézelin

INRA

Hélène Kiefer

UMR BDR, INRA, ENVA,
Université Paris Saclay

View Related Events