



## Comparative subproteomic analysis of experimentally induced fluoroquinolone resistance and ciprofloxacin stress in *Salmonella* Typhimurium DT104B

Susana Correia, Michel Hébraud, Ingrid Chafsey, Christophe Chambon, Didier Viala, Carmen Torres, Manuela Caniça, José-Luis Capelo, Patricia Poeta, Gilberto Igrejas

### ► To cite this version:

Susana Correia, Michel Hébraud, Ingrid Chafsey, Christophe Chambon, Didier Viala, et al.. Comparative subproteomic analysis of experimentally induced fluoroquinolone resistance and ciprofloxacin stress in *Salmonella* Typhimurium DT104B. 2. International Caparica Conference in Antibiotic Resistance (I-IC2AR), Jun 2017, Caparica, Portugal. 1 p., 2017. hal-02737755

**HAL Id: hal-02737755**

**<https://hal.inrae.fr/hal-02737755>**

Submitted on 2 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.




# Comparative subproteomic analysis of experimentally induced fluoroquinolone resistance and ciprofloxacin stress in *Salmonella* Typhimurium DT104B

Susana Correia<sup>1-4\*</sup>, Michel Hébraud<sup>5,6</sup>, Ingrid Chafsey<sup>5</sup>, Christophe Chambon<sup>6</sup>, Didier Viala<sup>6</sup>, Carmen Torres<sup>7</sup>, Manuela Caniça<sup>8</sup>, José Luis Capelo<sup>4,9</sup>, Patrícia Poeta<sup>3,4</sup> and Gilberto Igrejas<sup>1,2,4</sup>

<sup>1</sup>Functional Genomics and Proteomics Unit, University of Trás-os-Montes and Alto Douro, Portugal. <sup>2</sup>Department of Genetics and Biotechnology, University of Trás-os-Montes and Alto Douro, Portugal. <sup>3</sup>Veterinary Science Department, University of Trás-os-Montes and Alto Douro, Portugal. <sup>4</sup>UCIBIO-REQUIMTE, Faculty of Science and Technology, Nova University of Lisbon, Portugal. <sup>5</sup>Université Clermont Auvergne (UCA), Institut National de la Recherche Agronomique (INRA), UMR Microbiologie Environnement Digestif et Santé (MEDIS), site de Theix, France. <sup>6</sup>Institut National de la Recherche Agronomique (INRA), UR370 QuaPA, Plate-Forme d'Exploration du Métabolisme composante protéomique, site de Theix, France. <sup>7</sup>Área de Bioquímica y Biología Molecular, Universidad de La Rioja, Spain. <sup>8</sup>National Reference Laboratory of Antibiotic Resistances and Healthcare Associated Infections (NRL-AMR-HAI), Department of Infectious Diseases, National Health Institute Doutor Ricardo Jorge (INSA), Portugal. <sup>9</sup>ProteoMass Scientific Society, Faculty of Sciences and Technology, Campus de Caparica, Portugal. \*[scorreia@utad.pt](mailto:scorreia@utad.pt)


Fluoroquinolone-resistant *Salmonella* spp. is a global high priority pathogen for research and development of new antibiotics (World Health Organization, 2017) and *Salmonella* Typhimurium DT104B multiresistant strains with additional quinolone resistance have been responsible for global outbreaks and high mortality. Multiple factors are known to be involved in fluoroquinolone resistance; however, additional determinants continue to be described and the exact contribution of many mechanisms remains unexplained and/or need better understanding.

To give new insights into the mechanisms involved in fluoroquinolone resistance in *Salmonella* Typhimurium DT104B, this work aimed to evaluate subproteome changes between a laboratory-induced fluoroquinolone-resistant strain (Se6-M) and its parent strain (Se6) and also in Se6-M under ciprofloxacin (CIP) stress.



- A *Salmonella* Typhimurium DT104B isolate (Se6) was previously recovered from a hospitalized elderly patient with acute gastroenteritis before CIP treatment [1].
- An *in vitro* selected fluoroquinolone-resistant mutant (Se6-M) was experimentally obtained from the parental Se6 strain by a multi-step selecting process in the presence of increasing concentrations of CIP [2].
- Se6 and Se6-M were proven to be clonally related by PFGE.
- The different comparative subproteomic analyses were performed by combining 2-DE~LC-MSMS and shotgun LC-MSMS identification approaches for intracellular and membrane subproteomes, respectively.
- Proteomics data was validated by performing quantitative real time PCR for selected proteins of interest.

## SCHEMATIC REPRESENTATION OF THE MAIN RESISTANCE MECHANISMS IDENTIFIED



Periplasmic substrate-binding proteins of ABC importers:


ArtI, FliY, **GltI**/**GltI**, GsiB, LsrB, **MalE**, ModA, MppA, Oxd-6a, PstS, RbsB, **UgpB**, YrbC

Global regulators of bacterial metabolism and bacterial stress response proteins:

CspA, CspC, H-NS, IhfA, LexA, NusG, PhoP, RecA, Rho, SeqA, StpA,

Roles as determinants of antimicrobial resistance and/or targets for the development of new antimicrobial drugs

The proteomic signatures obtained provide valuable information about mechanism-related differential protein expression, supporting the current knowledge and leading to new testable hypotheses on the mechanisms of bacterial resistance to fluoroquinolones and other drugs.



Representative 2-DE gels with the protein spots identified by LC-MS/MS with significant differential abundance between the intracellular subproteomes of the *S. Typhimurium* DT104B strains (SameSpots software). Left: the protein spots more abundant in Se6 are represented on top and the more abundant in Se6-M are shown on the bottom. Right: the protein spots more abundant in Se6-M grown without ciprofloxacin stress are represented on top and the more abundant in Se20+CIP are shown on the bottom.

[1] de Toro M, et al (2010). *In vivo* selection of *aac(6')-Ib-cr* and mutations in the *gyrA* gene in a clinical *qnrS1*-positive *Salmonella enterica* serovar Typhimurium DT104B strain recovered after fluoroquinolone treatment. *J Antimicrob Chemother* 65(9): 1945-1949.

[2] Correia S, et al (2016). Impacts of experimentally induced and clinically acquired quinolone resistance on the membrane and intracellular subproteomes of *Salmonella* Typhimurium DT104B. *J Proteomics* 145: 46-59.