



HAL
open science

A model-based assessment of the soil C storage potential at the national scale: A case study from France

Sylvain Pellerin, Laure Bamière, Julie Constantin, Camille Launay, Raphaël Martin, Michele Schiavo, Denis Angers, Laurent Augusto, Jérôme Balesdent, Isabelle Basile-Doelsch, et al.

► **To cite this version:**

Sylvain Pellerin, Laure Bamière, Julie Constantin, Camille Launay, Raphaël Martin, et al.. A model-based assessment of the soil C storage potential at the national scale: A case study from France. Food security and climate change: 4 per 1000 initiative new tangible global challenges for the soil, Jun 2019, Poitiers, France. 10.15454/1.5433098269609653E12 . hal-02737947

HAL Id: hal-02737947

<https://hal.inrae.fr/hal-02737947>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License



A model-based assessment of the soil C storage potential at the national scale: A case study from France

Sylvain PELLERIN, Laure BAMIÈRE, Julie CONSTANTIN, Camille LAUNAY, Raphael MARTIN, Michele SCHIAVO, Denis ANGERS, Laurent AUGUSTO, Jérôme BALESSENT, Isabelle BASILE-DOELSCH, Valentin BELLASSEN, Rémi CARDINAEL, Lauric CECILLON, Eric CESCHIA, Claire CHENU, Joel DAROUSSIN, Philippe DELACOTE, Nathalie DELAME, François GASTAL, Anne-Isabelle GRAUX, Bertrand GUENET, Sabine HOUOT, Katja KLUMPP, Elodie LETORT, Manuel MARTIN, Bruno MARY, Safya MENASSERI, Delphine MEZIERE, Claire MOSNIER, Thierry MORVAN, Jean ROGER-ESTRADE, Laurent SAINT-ANDRE, Olivier THEROND, Valérie VIAUD, Olivier RECHAUCHERE, Guy RICHARD

Objectives

- To assess and map the potential for additional carbon storage in agricultural soils in mainland France following the implementation of soil C storing practices
- To assess their cost
- To propose an optimal cost-efficient strategy for additional soil C storage at the national scale



Method

Nine soil C storing practices were selected, based on a literature review

- no-tillage
- expansion of cover crops (longer and more frequent cover crops)
- new C inputs (not already spread on agricultural soils under current management practices)
- expansion of temporary grasslands (instead of silage maize)
- agroforestry
- hedges
- moderate intensification of extensive grasslands (+50kgN/ha)
- animal grazing instead of mowing
- grass cover of vineyard

For each selected practice, its potential applicability was calculated considering technical constraints (ex no reduced tillage for sugar beet, no cover crop for intercrop period < 2 months, no agroforestry if soil depth < 1m or plot size < 1ha...)


Method

The additional C storage following the implementation of C storing practices was assessed using a modelling approach at a fine spatial-scale resolution ($\approx 1 \text{ km}^2$)


Each grid cell was characterized by its:

- dominant land-use
- local climate
- dominant soil type
- dominant cropping system (crop rotation and current farming practices)
- initial soil C stock value


Land use




Climate



Clay content



Initial soil C stock value




Method

Two models were used for simulations

- STICS for arable crops
- PaSim for permanent grasslands

Both models include an explicit representation of the C cycle



Method

For each grid cell, changes in soil C stock were simulated for a period of 30 years


- under current management practices (= baseline)
- under C storing management practices (ex expansion of cover crops)

The additional C storage was calculated as the difference between the simulated soil C stock under C storing practices and the simulated soil C stock under current management practices

Simulations were performed under current climatic conditions

Simulations were performed for the 0-30cm horizon, and then extrapolated to the whole soil profile


For each C storing practice, a complete greenhouse gases budget was calculated





Example of results: expansion of cover crops

Percentage of intercrop periods with a cover crop

In the baseline (current farming practices)



In the scenario with expansion of cover crops




Potential applicability :
16.6 Mha
(96.1% of the cropland area)


Additional carbon storage provided by the expansion of cover crops

Average additional C storage

+126 ± 93 kg C/ha/yr



Map of the additional C storage provided by the expansion of cover crops



❖ Less additional carbon storage provided by cover crops in the western part of France;

- Due to crop rotations already including perennial temporary grasslands, which provide less opportunities for incorporating cover crops

❖ By contrast, more additional carbon storage in arable cropping systems of the Parisian Basin and South-western France

Aggregated results for all C-storing practices

	Additional C storage 0-30 cm soil layer	Potential applicability	Potential additional C storage at the national level 0-30 cm soil layer	Relative yearly increase of soil C stocks (=949 Mt C for cropland soils in mainland France)
	Kg C/ha/an	Mha	Mt C/year	‰ /year
Arable cropping systems				
Expansion of cover crops	+126	16.03	+2.019	
No tillage	+60	11.29	+0.677	
New carbon inputs	+61	4.21	+0.257	
Expansion of temporary grasslands	+114	6.63	+0.756	
Agroforestry	+207	5.33	+1.102	
Hedges	+17	8.83	+0.150	
Total for croplands			+4.960	+5.2 ‰

Aggregated results for all C-storing practices


	Additional C storage Horizon 0-30 cm	Potential applicability	Potential additional C storage at the national level Horizon 0-30 cm	Relative yearly increase of the soil C stock
	Kg C/ha/year	Mha	Mt C/year	‰ /year
Permanent grasslands				
Moderate intensification of extensive grasslands	+176	3.94	+0.694	
Grazing instead of mowing	+265	0.09	+0.023	
Total for permanent grasslands			+0.720	+0.9 ‰
Vineyard				
Grass cover	+182	0,56	+0.103	
Total for vineyard			+0.103	+3.7 ‰
Total for French agricultural soils (without forests)			5,78	+3,3 ‰

- A potential for additional C storage of about 5.78 Mt C/year (in the 0-30cm soil layer)
- This represents an annual increase of +3.3‰ for agricultural soils
- Extrapolated to the whole soil profile (5.78 → 8.43 MtC = 31MtCO₂e), this additional C storage would compensate
 - 6.8% of national GHG emissions (458 MtCO₂e)
 - 41% of emissions from the French agricultural sector (76.7 MtCO₂e)


Costs

C storing practice	Potential applicability (Mha)	Cost for farmer (€/ha/year)	Storage cost (€/tC)	Storage cost (€/tCO ₂)
New carbon inputs	4,21	-52 (-117; -8)	-494 (-1 192; -134)	-135 (-325; -37)
Grass cover of vineyards permanent	0,15	-26 (-27; -22)	-56 (-44; -77)	-15 (-21; -11)
In winter	0,41	-15 (15; 15)	-51 (-51)	-14 (-14)
Expansion of cover crops	16,03	39 (12; 147)	180 (69; 1 104)	49 (19; 301)
Moderate intensification of extensive grasslands	3,94	28 (12; 38)	130 (60, 1 189)	35 (16;324)
Grazing instead of mowing	0,09	73 (-85; 146)	203 (-2 791; 518)	55 (-761; 141)
Expansion of temporary grasslands	6,63	91 (-40; 263)	473 (-242; 1 667)	129 (-66; 455)
Agroforestry	5,33	118 (63; 179)	302 (195; 386)	82 (53; 105)
Hedges	8,83	73 (54; 87)	2 322 (2 013; 3 618)	633 (549; 987)

Marginal carbon storage cost curve



Contributing C storage practices at the shadow price of carbon (201,7 €/tC = 55 €/tCO₂e)



❖ Additional C storage : 4,06 MtC/year

❖ Total cost= 159 M€/year

❖ Contributing C storing practices :

- Cover crops : 58%
- Moderate intensification of extensive grasslands : 19%
- New C inputs : 11%
- Temporary grasslands : 6%

Summary

- A potential for additional C storage of about 5,78 Mt C/year (in the 0-30cm soil layer)
- This represents a relative increase of +3,3‰ for agricultural soils (+ 5,2‰ if only arable soils are considered)
- This potential is mainly found in arable soils (86% of the total potential), partly because initial soil C stocks are low
- Extrapolated to the whole soil profile, this additional C storage would compensate
 - 6,8% of national GHG emissions
 - 41% of emissions from the agricultural sector
- About half of this potential is at a lower cost than the shadow price of carbon