



HAL
open science

Mycotoxin biosynthesis and central metabolism are two interlinked pathways in *Fusarium graminearum*, as demonstrated by the extensive metabolic changes induced by caffeic acid exposure

Sylvain Chéreau, Vessela Atanasova-Penichon, Laurie Legoahec, Stéphane Bernillon, Catherine Deborde, Mickael Maucourt, Marie-Noëlle Verdal-Bonnin, Laetitia Pinson-Gadais, Annick Moing, Nadia Ponts, et al.

► **To cite this version:**

Sylvain Chéreau, Vessela Atanasova-Penichon, Laurie Legoahec, Stéphane Bernillon, Catherine Deborde, et al.. Mycotoxin biosynthesis and central metabolism are two interlinked pathways in *Fusarium graminearum*, as demonstrated by the extensive metabolic changes induced by caffeic acid exposure. 14. European Fusarium Seminar, Universität für Bodenkultur Wien [Vienne, Autriche] (BOKU). AUT., Apr 2018, Tulln, Austria. 122 p. hal-02737991

HAL Id: hal-02737991

<https://hal.inrae.fr/hal-02737991>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



April 8 – 11, 2018
Minoritenkloster
Tulln, AUSTRIA

ORGANIZING COMMITTEE

Gerhard Adam
(BOKU, AT)

Rainer Schuhmacher
(BOKU, AT)

Hermann Bürstmayr
(BOKU, AT)

Joseph Strauss
(BOKU, AT)

Rudolf Krska
(BOKU, AT)

Gerlinde Wiesenberger
(BOKU, AT)

Dieter Moll
(BIOMIN, AT)

Alexandra Khassidov
(ÖGMBT, AT)

INTERNATIONAL SCIENTIFIC COMMITTEE

**Sarah M. De
Saeger**
(BE)

Ruth Dill-Macky
(US)

Doris Marko
(AT)

**Thomas
Miedaner**
(DE)

Antonio Moretti
(IT)

Michele Suman
(IT)

B. Gillian Turgeon
(US)

Table of Contents

Table of Contents	2
General Information	3
Welcome Address	4
Invited Speakers	5
Detailed Program	13
SUNDAY - April 8	13
MONDAY - April 9	13
TUESDAY - April 10	16
WEDNESDAY - April 11	19
Posters Table	21
Lectures	22
Welcome Note & Plenary Session	23
Session 1: Fusarium genomics and virulence mechanisms, population genetics and diversity	25
Session 2: Host resistance: mechanisms, genetics, genomics and breeding	32
Session 3: Fusarium secondary metabolites and metabolomics of Fusarium-host plant interactions	39
Session 4: Fusarium mycotoxins - Toxicology, Metabolism and Remediation	46
Session 5: Integrated Fusarium management (pre/postharvest, epidemiology and forecasting, fungicide resistance, utilization of contaminated batches)	54
Poster Presentations	61
Author Index	110
Participant Index	115
Imprint	123

MB-03 Metabolomics of Fusarium head blight: Examining the attack of Fusarium graminearum during infection of two near isogenic wheat lines differing in the resistance QTLs Fhb1 and Qfhs.ifa-5A

Alexandra Simader¹, Sven Gottwald², Bernhard Wolf¹, Christoph Büschl¹, Maria Doppler¹, Barbara Steiner², Hermann Bürstmayr², Gerhard Adam³, Rainer Schuhmacher¹

¹ Department of Agrobiotechnology, IFA-Tulln, Center for Analytical Chemistry, Austria

² Institute for Biotechnology in Plant Production, Department of Agrobiotechnology (IFA-Tulln), University of Natural Resources and Life Sciences, Vienna (BOKU), Tulln, Austria

³ Institute of Applied Genetics and Cell Biology, University of Natural Resources and Life Sciences, Vienna (BOKU), Tulln, Austria

The fungus *Fusarium graminearum* (*Fg*) is able to infect wheat and other small grain cereals. Major problems arising from this so called Fusarium head blight (FHB) disease comprise yield and quality losses as well as mycotoxin contamination of the affected crop. In the present study a greenhouse experiment has been carried out with the aim to investigate fungal attack of two near isogenic wheat lines differing in the FHB resistance QTLs Fhb1 and Qfhs.ifa-5A. To this end, we have developed an isotope-assisted, untargeted metabolomics approach to distinguish fungal metabolites from those produced by the affected plants.

Within a time course experiment encompassing 3, 6, 12, 24, 36, 48, 72 and 96 hours after infection (hai), about 100 metabolites were determined to be explicitly deriving from *Fg*. For another approximately 250 metabolites, all of which were found to be induced by fungal infection, we were able to classify as being either produced by *Fg* to support infection, the plant in defence or by both interaction partners. Ca. 50% of the fungal metabolites could be annotated including those belonging to the type B trichothecene pathway. Many of the *Fg* assigned metabolites were found to be produced in both genotypes. Towards the end of the time course, most of the detected fungal metabolites were more abundant in the susceptible wheat genotype, lacking the two resistance QTLs. Interestingly, at an earlier stage during infection (about 48 hai) some of these fungal compounds were found to be more abundant in the resistant compared to the susceptible genotype. Our data suggest that in the resistant cultivar, the fungus is either “forced” to produce more and higher amounts of secondary metabolites at an early infection stage or alternatively, higher toxin formation by *Fg* in the resistant wheat genotype may help the plant to induce a quicker/stronger response against the pathogen including the detoxification of mycotoxins via glycosylation. Towards the end of the monitored time window (96 hai) we observed that *Fg* is not only able to produce higher amounts of well-known toxins, but also to produce a much higher number of metabolites in the infected plants, as was particularly observed for the susceptible wheat line.

MB-04 Mycotoxin biosynthesis and central metabolism are two interlinked pathways in *Fusarium graminearum*, as demonstrated by the extensive metabolic changes induced by caffeic acid exposure

Sylvain Chéreau¹, Vessela Atanasova-Pénichon¹, Laurie Legoahec¹, Stéphane Bernillon², Catherine Deborde², Mickaël Maucourt², Marie-Noëlle Verdal-Bonnin¹, Laëtitia Pinson-Gadai¹, Annick Moing², Nadia Ponts¹, Florence Richard-Forget¹

¹ UR1264 MycSA, INRA, France

² Metabolome Facility of Bordeaux Functional Genomics Center, IBVM, Centre INRA Nouvelle Aquitaine Bordeaux, 71 Avenue Edouard Bourlaux, 33140 Villenave d'Ornon, France

Fusarium graminearum is a major plant pathogen that causes Fusarium Head Blight in wheat and Gibberella Ear Rot in maize and produces type B trichothecene mycotoxins (TCTB) in infected grains. A comprehensive understanding of the molecular and biochemical mechanisms underlying the regulation of TCTB biosynthesis is required for improving or developing novel strategies to efficiently manage the risks posed by *F. graminearum*. Elucidation of the association of TCTB biosynthesis with other metabolic processes, central and specialized ones was the focus of this study. Combined ¹H-NMR and LC-QTOF-MS analyses were used to investigate and compare the exo- and endo- metabolomes of *F. graminearum* grown in toxin-inducing and repressing caffeic acid-conditions. Ninety-five metabolites were putatively or unambiguously identified including 26 primary and 69 specialized metabolites. Our data demonstrated that the inhibition of TCTB production induced by caffeic acid exposure was associated with significant changes in secondary and primary metabolism of *F. graminearum* although the fungal growth was not affected. The main metabolic changes were an increase in the accumulation of several polyketides including toxic ones, alterations in the tricarboxylic organic acid cycle and modifications in the metabolisms of some amino-acids and sugars. While these findings allow improving the knowledge on the mechanisms that govern the inhibition of TCTB production by caffeic acid, they also demonstrate the interdependence between the biosynthetic pathway of TCTB and several additional metabolic pathways including primary and specialized ones. These outcomes provide further evidence of the multifaceted role of TCTB in the life cycle of *F. graminearum*.