

The effect of perennial rygrass cultivars and allowance on utilisation, grazing efficiency and milk production

J. Mcdonagh, T.J. Gilliland, Mary Mcevoy, Luc Delaby, M. O'Donovan

► **To cite this version:**

J. Mcdonagh, T.J. Gilliland, Mary Mcevoy, Luc Delaby, M. O'Donovan. The effect of perennial rygrass cultivars and allowance on utilisation, grazing efficiency and milk production. 26. General meeting of the European Grassland Federation (EGF), Sep 2016, Trondheim, Norway. hal-02738593

HAL Id: hal-02738593

<https://hal.inrae.fr/hal-02738593>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The effect of perennial ryegrass cultivars and allowance on utilisation, grazing efficiency and milk production

McDonagh J.^{1,2}, Gilliland T.J.^{2,3}, McEvoy M.¹, Delaby L.⁴ and O'Donovan M.¹

¹Animal & Grassland Research and Innovation Centre, Teagasc, Moorepark, Co Cork; ²Institute for Global Food Security, School of Biological Sciences, Queen's University, Belfast, N. Ireland; ³Agrifood & Biosciences Institute, Hillsborough, BT26 5DR, N. Ireland; ⁴Institut National de Recherche Agronomique, Agrocampus Ouest, UMR 1348, Physiologie, Environnement et Génétique Pour l'Animal et les Systèmes d'Élevage, 35590 Saint Gilles, France; michael.odonovan@teagasc.ie

Abstract

The objective of this study was to investigate the effect of four cultivars (AstonEnergy, Delphin, Glenroyal and Tyrella) on sward utilisation, milk production and grazing efficiency of lactating dairy cows at two daily herbage allowances (DHA) during the vegetative growth phase. Four cultivars were compared by offering cows two levels of DHA 15 and 20 kg dry matter (DM) cow⁻¹ day⁻¹ (>4cm), resulting in 8 treatments. Increasing herbage allowance from 15 to 20 kg DM cow⁻¹ day⁻¹ increased milk yield and solids, but reduced sward utilisation with grazing efficiency being unaffected. AstonEnergy had a longer free leaf lamina ($P < 0.001$; + 4.6 cm) but also higher leaf ($P < 0.001$; + 0.14) and lower stem ($P < 0.01$; -0.08) than Glenroyal. This resulted in a higher sward utilisation ($P < 0.001$; +12%) milk yield ($P < 0.05$; +1 kg) and milk solids ($P < 0.05$; +0.1 kg) than Glenroyal. Delphin was most similar to AstonEnergy in sward structure and gave similar sward utilisation and milk production, whereas Tyrella was broadly equivalent to an average of the examined cultivars and also supported an intermediate milk output. Cultivars with longer free-leaf-lamina and higher leaf content were associated with reduced number of grazing bites ($P < 0.05$) and ruminating time ($P < 0.001$).

Keywords: ryegrass, grazing, milk production, sward utilization

Introduction

Improved efficiency of animal production from grass is the ultimate goal of forage grass breeding for most temperate regions (Wilkins and Humphreys, 2003). Perennial ryegrass cultivars form the basis of pasture production in many temperate regions. Sward structure can vary between cultivars and has been shown to affect milk production of grazing dairy cows (Wims *et al.*, 2013), though a significant effect on milk production does not always occur (Tas *et al.*, 2006). In addition to affecting milk production, sward structure is also an important determinant of grazing efficiency (O'Donovan *et al.*, 2004) even more, where cows are managed on intensive intervals of rotational grazing which require rapid removal of herbage. Confirmation and quantification of animal responses to sward structural differences is a critical step in comparing perennial ryegrass cultivars for on farm performance. The objective of this study was to compare the milk production performance of dairy cows when grazing four perennial ryegrass cultivars varying in sward structure at two herbage allowances.

Materials and methods

Four perennial ryegrass cultivars comprising two diploids (D) and two tetraploids (T) were sown as monocultures. Cultivars included AstonEnergy (T), Delphin (T), Glenroyal (D) and Tyrella (D). Seven cows (two primiparous and five multiparous) were assigned to each cultivar during the vegetative growth phase (1 July to 28 July) at two herbage allowances: 15 and 20 kg dry matter (DM) cow⁻¹ day⁻¹. This resulted in 8 treatments (4 cultivars × 2 allowances). Cows were blocked based on lactation number (3.0), milk yield (18.5 kg d⁻¹), milk solids (1.4 kg d⁻¹), protein (33.7 g kg⁻¹), fat (40.5 g kg⁻¹) and lactose (47.6 g kg⁻¹), and body weight (549 kg) and body condition score (3.0; based on a scale of 1 to 5, 1 = emaciated,

5 = extremely fat). Fresh grass was allocated to cows daily. Pre- and post-grazing sward heights were measured daily (50 heights per treatment) directly before and after grazing using a rising plate meter with a steel plate (diameter 355 mm, 3.5 kg m⁻²; Jenquip, Fielding, New Zealand). Pre-grazing free-leaf-lamina (FLL) was measured from the highest ligula (longest leaf sheath) to the top of the leaf using a hand-ruler on 100 random tillers across each offered treatment twice weekly. Approx. 40 g of representative herbage was sampled from each treatment twice weekly to ground level; samples were then cut to the weekly post-grazing height per respective treatment and separated manually into leaf and stem fractions. Fractions were then dried for 16 h at 90 °C for DM determination. Grazing behaviour was estimated by fitting all cows simultaneously with an Institute of Grassland and Environmental Research (IGER) headset behaviour recorder after morning milking and removed prior to the morning milking 24 hours later. Individual milk yields were recorded at individual AM and PM milking. Milk solids (protein, fat and lactose) were measured twice weekly. Animal variables were analysed using PROC MIXED in SAS (2011), with grass allowance, cultivar, their interactions and pre-experiment covariate included in the model as fixed effects and animal included as a random effect.

Results and discussion

There was no cultivar × allowance interaction for any of the sward, animal or grazing efficiency variables in this study. Cows allocated 15 kg DM had a higher pseudostem proportion ($P < 0.05$; +0.2) in the grazed sward compared to the 20 kg DM daily herbage allowances (DHA) grazed swards. Herbage allowance of 20 kg DM resulted in a lower utilisation ($P < 0.001$; -23%), higher post-grazing height, longer post-grazing FLL and SH ($P < 0.001$; +1.1 cm, +1.5 cm, +0.9 cm, +0.7 cm) of the grazed swards in comparison to the 15 kg DM DHA swards (Table 1). As a result, cows on the higher DHA produced significantly more milk yield and milk solids ($P < 0.001$; +1.6, +1.1 kg day⁻¹). Although the four varieties had a similar herbage mass (kg DM above 4 cm from 12 m², not shown), AstonEnergy and Delphin were grazed to a lower residual height (4.1, 4.2 cm) compared to the highest, Glenroyal (+0.6 cm; $P < 0.001$). AstonEnergy also had higher FLL ($P < 0.001$), higher leaf ($P < 0.001$) and lower stem ($P < 0.05$) contents than Glenroyal. Importantly, AstonEnergy supported a higher milk yield (+1 kg cow⁻¹ d⁻¹; $P < 0.05$) and milk solids (+0.1 kg cow⁻¹ d⁻¹; $P < 0.05$) than Glenroyal. Delphin was most similar to AstonEnergy in sward structure and gave the same milk production, whereas Tyrella was broadly equivalent to an average of the examined cultivars and also supported an intermediate milk output. This is in agreement with O'Donovan and Delaby (2005) stating that cultivars producing a higher green leaf allowance impact positively on milk production. Cows grazing AstonEnergy with a longer FLL and higher leaf:stem ratio had significantly less grazing bites ($P < 0.001$; -7083) than cows grazing Glenroyal which had increased ruminating mastications ($P < 0.001$; +6717) compared to the other three cultivars averaging 32,329 (Figure 1). Higher stem content resulted in failure to break the barrier of standing stem despite elevated

Table 1. The effect of perennial ryegrass cultivars and herbage allowance on milk production and solids, post-grazing sward height, pre-grazing free-leaf-lamina, leaf and stem proportion and sward utilisation.

Herbage DM allowance	15 kg	20 kg	Aston ¹	Delphin	Glenroyal	Tyrella	SED	Allowance	Cultivar
Milk yield (kg cow ⁻¹ d ⁻¹)	15.1 ^a	16.7 ^b	16.3 ^a	16.3 ^a	15.3 ^b	15.7 ^{ab}	0.26	***	*
Milk solids (kg cow ⁻¹ d ⁻¹)	1.24 ^a	1.35 ^b	1.31 ^a	1.35 ^a	1.23 ^b	1.28 ^{ab}	0.03	***	*
Post-grazing SH (cm)	3.8 ^a	4.9 ^b	4.1 ^a	4.2 ^a	4.7 ^c	4.4 ^b	0.19	***	***
Pre-grazing FLL (cm)	17.5	17.5	18.7 ^{ab}	19.9 ^a	14.1 ^c	17.4 ^b	0.37	NS	***
Leaf (%)	74.0	77.0	82.0 ^a	78.5 ^{ab}	68.0 ^c	74.0 ^b	0.01	NS	***
Stem (%)	17.0	15.0	12.0 ^a	15.5 ^{ab}	19.5 ^b	16.0 ^b	0.01	NS	*
Utilisation (%)	105 ^a	82 ^b	99 ^a	96 ^a	87 ^c	92 ^b	0.01	***	***

¹ Aston = AstonEnergy.

^{a-c} Means within a row with different superscripts differ ($P < 0.05$).




Figure1. The effect of cultivar on grazing bites and ruminating mastications of grazing dairy cows.

grazing pressure (Parga *et al.*, 2000) and impacted negatively on rumination process (Van Soest *et al.*, 1991) in turn reducing milk production.

Conclusions

To conclude, increased herbage allowance increases milk production but reduces sward utilisation significantly. Variations in sward structure in particular free leaf lamina between grass cultivars influence sward utilisation. When sward structural characteristics such as leaf : stem ratio and free leaf lamina length coupled with digestibility are differing by the magnitudes observed in this study between the four cultivars, milk production and grazing efficiency of grazing dairy cows are influenced. Improving cultivar sward characteristics such as leaf:stem ratio and free leaf lamina have the potential to improve overall efficiency of the conversion of grass to milk and improve grazing efficiency.

References

- O'Donovan M. and Delaby L. (2005) A comparison of perennial ryegrass cultivars differing in heading date and grass ploidy with spring calving dairy cows grazed at two different stocking rates. *Animal Research* 54, 337-350.
- O'Donovan M., Delaby L. and Peyraud J.L. (2004) Effect of time of initial grazing date and subsequent stocking rate on pasture production and dairy cow performance. *Animal Research* 53, 489-502.
- Parga J., Peyraud J.L. and Delagarde R. (2000) Effect of sward structure and herbage allowance on herbage intake and digestion by strip-grazing dairy cows. In: Rook A.J. and Penning P.D. (eds). *Occasional Symposium No. 34*. British Grassland Society, pp. 61-66.
- Tas, B.M., Taweel S., Smit H.Z., Elgersma A., Dijkstra J. and Tamminga, S. (2006) Effects of Perennial Ryegrass Cultivars on Intake, Digestibility, and Milk Yield in Dairy Cows. *J. Dairy Sci.* 89, 3494-3500.
- Van Soest P.J, Robertson J.B. and Lewis B.A. (1991) Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *Journal of Dairy Science* 74, 3583-3597.
- SAS (2011) *Version 9.3*. SAS Institute. Cary, NC, USA.
- Wilkins P.W. and Humphreys M.O. (2003) Progress in breeding perennial forage grasses for temperate agriculture. *The Journal of Agricultural Science* 140, 129-150.
- Wims C.M., McEvoy M., Delaby L., Boland T.M. and O'Donovan M. (2013) Effect of perennial ryegrass (*Lolium perenne* L.) cultivars on the milk yield of grazing dairy cows. *Animal* 7, 410-421.