

Identification of uterine ionic transport proteins involved in providing the mineral precursors for eggshell formation in hens

Aurélien Brionne, Yves Y. Nys, Christelle Hennequet-Antier, Joël Gautron

▶ **To cite this version:**

Aurélien Brionne, Yves Y. Nys, Christelle Hennequet-Antier, Joël Gautron. Identification of uterine ionic transport proteins involved in providing the mineral precursors for eggshell formation in hens. 16. Journées françaises de biologie des tissus minéralisés, May 2014, Limoges, France. hal-02738600

HAL Id: hal-02738600
https://hal.inrae.fr/hal-02738600

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



FICHE DE SOUMISSION de RESUME POUR LES JFBTM16

Du 14 au16 Mai 2014 à Limoges

NB Cette fiche comportant 2 pages est à compléter en plus de la procédure d'inscription en ligne sur le site de la SFBTM et à renvoyer par email (reine.bareille@inserm.fr), **avant le 31 Mars 2014*** en mettant comme nom du fichier: le nom de l'orateur suivi de jfbtm15 au format .doc (ex: Dupont-jfbtm16.doc).
Si le même auteur soumet un deuxième résumé, faire une seconde fiche (ex: Dupont2-jfbtm16.doc)

Nom, **Prénom** : BRIONNE Aurélien

Adresse professionnelle complète : INRA Centre VAL DE LOIRE, UR83 Recherches Avicoles, F-37380 Nouzilly, France.
Adresse email : aurelien.brionne@tours.inra.fr
Tél. : 02 47 42 76 87
Fax :

- **Thème de recherche** (indiquer le n° choisi sur la liste) : 2

1 - Biologie cellulaire et Moléculaire. 2 - Biominéralisation.
3 - Développement et différenciation. 4 – Génétique et Physiologie.
5 - Ostéoimmunologie 6 – Biomatériaux

- **Le travail concourt au prix des jeunes chercheurs (Master/Thèse)**: **OUI** **NON**

(NB joindre un scan de la photocopie de la carte d'étudiant (format: Dupont-étudiant.pdf) ; l'étudiant qui concourt pour une bourse doit être le présentateur et apparaitre comme premier auteur du résumé)

- **Type de Présentation souhaitée** : **Orale** **Affichée**

(Ce choix sera validé par le comité scientifique et vous sera communiqué Mi-Avril 2014)

- **Présentation : Etudiant (Master/Doctorant)** **Post-Doctorant** **Statutaire**

- **Proposer 4 mots clés:** **1- chicken**
  2- eggshell
  3- uterus
  4- ionic transport

NB : Afin d'éviter une édition lourde des résumés pour les homogénéiser, merci d'avance de respecter les consignes de formatage indiquées sur la page suivante.
**Après le 31 Mars 2014, les résumés seront acceptés en communication affichée*



Identification of uterine ionic transport proteins involved in providing the mineral precursors for eggshell formation in hens.

BRIONNE Aurélien[1], NYS Yves[1], HENNEQUET-ANTIER Christelle[1], and GAUTRON Joël[1]

[1]INRA, UR83 Recherches Avicoles, F-37380 Nouzilly, France

The bird eggshell is a complex bioceramic formed in the uterus segment of the chicken oviduct. The shell is made of 95% calcium carbonate (calcite) and 3.5% organic matrix made of proteins, glycoproteins and proteoglycans. The prerequisite for shell mineralization is the supply of large amounts of calcium and bicarbonate in the uterine fluid where calcification takes place. Both ions, are supplied by the blood via trans-epithelial transport in the uterus, and require ion channels, ion pumps and ion exchangers. The present study describes and updates the general model for ion transfers across the uterine tubular gland cells during eggshell formation. Our study used transcriptomic approach to reveal genes highly expressed in laying hen uterus during the active phase of mineralization, compared to the same period and tissue but following suppression of shell calcification by premature expulsion of the egg. This suppression abolishes the uterine egg mechanical stimulation and the calcium demand for shell formation. From the total list of genes over-expressed in uterus during eggshell calcification, we have identified 16 proteins related to ion transport updating the earlier model of ionic transfer.

Calcium is not stored in uterus but is continuously supplied from the blood. Hen uterine glandular cells transfer large amounts of calcium into the uterine lumen against the concentration gradient while preserving a low level of intracellular calcium. We observed that calbindin D28K was 17 times over-expressed during calcification. Low free calcium levels in the cell are maintained by calcium uptake by the endoplasmic reticulum via ATP depending calcium pumps as suggested by a 3 fold over-expression of ATP2A3 when the egg is being calcified. Finally, the calcium pumps ATP2B1 and ATP2B2, which can extrude calcium from the cytosol to an extracellular fluid against a strong electrochemical gradient, were 2 times over-expressed in our study, suggesting their involvement in the uterine Ca secretion. Carbonate of the eggshell is not derived from the blood HCO_3^- but rather from the plasma CO_2, which is hydrated in the uterus to produce bicarbonate. Carbonic anhydrase 2 is over-expressed at a very high level during calcification and should play a pivotal role for conversion of intracellular CO_2 to HCO_3^- in hen uterus.

The transfer of calcium and bicarbonate from the blood to the uterine fluid involves additional ion transfers o maintain cell homeostasis. We reported here transporters involved in Na^+, K^+, Mg^{2+}, H^+, and Cl^- exchanges to maintain physiological ionic concentrations in the cell.

This experimental approach has been used to reveal new candidate proteins and enrich the model describing ion transfers through the glandular cells of the uterus. Additional studies are still needed to quantify the respective role of ionic transporters which are controlling shell formation.