



HAL
open science

Deletion of the C-terminal part of helix alpha 2 does not prevent the prion conversion

Carola Munoz Montesino, Mohammed M. Moudjou, Laetitia Herzog, Danica Ciric, Human Rezaei, Vincent Béringue, Christina Sizun, Michel Dron

► To cite this version:

Carola Munoz Montesino, Mohammed M. Moudjou, Laetitia Herzog, Danica Ciric, Human Rezaei, et al.. Deletion of the C-terminal part of helix alpha 2 does not prevent the prion conversion. International Prion Congress 2014, May 2014, Trieste, Italy. 10.4161/pri.29369 . hal-02738722

HAL Id: hal-02738722

<https://hal.inrae.fr/hal-02738722>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

This article was downloaded by: [138.102.135.242]

On: 12 March 2015, At: 02:28

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Prion

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/kprn20>

Protein Structure and Biology

Published online: 01 Apr 2014.



[Click for updates](#)

To cite this article: (2014) Protein Structure and Biology, Prion, 8:sup1, 25-58, DOI: [10.4161/pri.29369](https://doi.org/10.4161/pri.29369)

To link to this article: <http://dx.doi.org/10.4161/pri.29369>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Versions of published Taylor & Francis and Routledge Open articles and Taylor & Francis and Routledge Open Select articles posted to institutional or subject repositories or any other third-party website are without warranty from Taylor & Francis of any kind, either expressed or implied, including, but not limited to, warranties of merchantability, fitness for a particular purpose, or non-infringement. Any opinions and views expressed in this article are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor & Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

It is essential that you check the license status of any given Open and Open Select article to confirm conditions of access and use.

P.42: Deletion of the C-terminal part of helix alpha 2 does not prevent the prion conversion

Carola Munoz-Montesino¹ Mohammed Moudjou,¹
Laetitia Herzog,¹ Daniča Ciric,¹ Human Rezaei,¹
Vincent Béringue,¹ Christina Sizun,² and Michel Dron¹

¹INRA; UR892 Virologie Immunologie Moléculaires; Jouy-en-Josas, France; ²CNRS; UPR2301 Institut de Chimie des Substances Naturelles; Gif-sur-Yvette, France

The mechanistic insights of prion conversion remain highly controversial. Several regions of PrP have been postulated as critical in this process; however, few attempts have been done to determine to what extent we can modify PrP sequences maintaining its properties. Recently, our laboratory progressed on these issues by showing that a PrP with eight extra amino acids inserted in the middle of the helix-2-helix-3 domain remains convertible into prion. These insertion mutants displayed no significant loss of the alpha helix content suggesting a replacement by the amino acids inserted. To determine whether a full size of H2 is required for prion conversion we performed series of deletions by site-directed mutagenesis in the ovine PrP sequence (VRQ haplotype). Mutants of PrP were then stably transfected in RK13 cells where protein expression and distribution of the mutants were monitored. The secondary structure of the recombinant proteins with similar deletions was analyzed by circular dichroism, showing no significant differences. The efficiency of infection was then tested through determination of the amount of protease resistant PrP accumulated in cells several passages after infection. Using 3 different prion strains for infection, we demonstrated that deletion of two turns of the H2 end does not prevent prion conversion indicating that this portion of the protein is not critically involved in the conformational change. We also probed that the susceptibility to infection was not altered in the mutants when compared to the wild type protein. The cells did propagate bona fide prions, infectious not only for naive homologous cell cultures but also for wt-expressing cells and for tg338 mice expressing the ovine PrP. This work demonstrates that the end of H2 is dispensable for prions and extends our previous conclusions that the amino acid specificity of this region was not required for the conversion.

To our best knowledge, most deletions in the protease-resistant domain that were reported so far remained unconvertible into prions with the exception of PrP106 produced by two large deletions in the first moiety of PrP^C. Our work demonstrates the possibility to produce prions with short internal deletions inside the protease-resistant core.

P.43: Cloning the prion and the Shadoo proteins: Restriction enzyme Body Doubles and PCR cloning

Eszter Tóth¹ Krisztina Huszár,² Petra G Bencsura,²
Péter István Kulcsár,¹ Barbara Vodicska,¹ Antal Nyeste,¹
Zsombor Welker,² Szilvia Tóth,² and Ervin Welker^{1,2}

¹Institute of Biochemistry; Biological Research Centre; Hungarian Academy of Sciences; Szeged, Hungary; ²Institute of Molecular Pharmacology; Research Centre for Natural Sciences; Hungarian Academy of Sciences; Budapest, Hungary

Here we demonstrate a procedure that solves a frequent problem in genetic engineering experiments that we faced working with the cDNA of the Prion and Shadoo proteins. This procedure allows the cloning of PCR fragments containing a recognition site of the restriction endonuclease (Type IIP) used for cloning in the sequence of the insert. A Type IIS endonuclease - a Body Double of the Type IIP enzyme - is used to generate the same protruding palindrome. Thus, the insert can be cloned to the Type IIP site of the vector without digesting the PCR product with the same Type IIP enzyme. We achieve this by incorporating the recognition site of a Type IIS restriction enzyme that cleaves the DNA outside of its recognition site in the PCR primer in such a way that the cutting positions straddle the desired overhang sequence. Digestion of the PCR product by the Body Double generates the required overhang. Hitherto the use of Type IIS restriction enzymes in cloning reactions has only been used for special applications, the approach presented here makes Type IIS enzymes as useful as Type IIP enzymes for general cloning purposes. To assist in finding Body Double enzymes, we summarised the available Type IIS enzymes which are potentially useful for Body Double cloning and created an online program (http://group.szbk.u-szeged.hu/welkergr/body_double/index.html) for the selection of suitable Body Double enzymes and the design of the appropriate primers. We routinely use this procedure to bypass this frequently arising problem.

P.44: Conformational properties of prion strains can be transmitted to recombinant prion protein fibrils in real-time quaking-induced conversion

Kazunori Sano, Ryuichiro Atarashi, and Noriyuki Nishida

Department of Molecular Microbiology and Immunology; Nagasaki University
Graduate School of Biomedical Sciences; Nagasaki, Japan

The phenomenon of prion strains with distinct biological characteristics has been hypothesized to be involved in the diverse structures of abnormal prion protein (PrP^{Sc}). However, the molecular basis of the strain diversity, including how to transmit the strain properties, remains uncertain. Real-time quaking-induced conversion (RT-QUIC) is a cell-free system that uses *E. coli*-derived recombinant PrP (rPrP) for the sensitive detection of PrP^{Sc}. To investigate whether properties of various prion strains can be transmitted to amyloid fibrils consisting of