Image analysis methods for assessment of H₂O₂ production and *Plasmopara viticola* development in grapevine leaves: Application to the evaluation of resistance to downy mildew

Ian Li Kim Khiook, Charles Schneider, Marie-Claire Héloir, Benjamin Bois, Xavier Daire, Marielle Adrian, Sophie Trouvelot

► To cite this version:

Ian Li Kim Khiook, Charles Schneider, Marie-Claire Héloir, Benjamin Bois, Xavier Daire, et al.. Image analysis methods for assessment of H₂O₂ production and *Plasmopara viticola* development in grapevine leaves: Application to the evaluation of resistance to downy mildew. 16. International Congress on Molecular Plant-Microbe Interactions (XVI IS-MPMI), Jul 2014, Rhodes, Greece. 2014. hal-02738757

HAL Id: hal-02738757

<https://hal.inrae.fr/hal-02738757>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

P195

IMAGE ANALYSIS METHODS FOR ASSESSMENT OF H₂O₂ PRODUCTION AND *PLASMOPARA VITICOLA* DEVELOPMENT IN GRAPEVINE LEAVES: APPLICATION TO THE EVALUATION OF RESISTANCE TO DOWNY MILDEW

Ian Li Kim Khiook¹, Charles Schneider¹, Marie-Claire Heloir¹, Benjamin Bois², Xavier Daire¹, Marielle Adrian¹, Sophie Trouvelot¹

¹Université de Bourgogne, UMR AgroSup/INRA/uB 1347 Agroécologie, ERL CNRS 6300, BP 86510, 21000 Dijon, France, ²Institut Universitaire de la Vigne et du Vin — CRC, UMR 6282 Biogéosciences CNRS/Université de Bourgogne-6, Boulevard Gabriel, 21000 Dijon, France

The grapevine downy mildew (*Plasmopara viticola*) provokes severe damages and destroys the harvest in the absence of an effective protection. Numerous fungicide treatments are thus generally necessary. To promote a sustainable production, alternative strategies of protection including new antifungal molecules, resistant genotypes or elicitor-induced resistance are under trial. To evaluate the relevance of these strategies, resistance tests are required. In this context, three image analysis methods were developed to read the results of tests performed to assess *P. viticola* sporulation and mycelial development, and H₂O₂ production in leaves. They have been validated using elicitors of plant defenses. These methods are reliable, innovative, rapid, and their modular concept allows their further adaptation to other host-pathogen systems.