



HAL
open science

Toward the development of functional foods for elderly

Aurélie Promeyrat, Thierry T. Astruc, Annie A. Venien, Claude C. de Oliveira Ferreira, Véronique Santé-Lhoutellier

► To cite this version:

Aurélie Promeyrat, Thierry T. Astruc, Annie A. Venien, Claude C. de Oliveira Ferreira, Véronique Santé-Lhoutellier. Toward the development of functional foods for elderly. 28. EFFoST Conference "Innovations in Attractive and Sustainable Food for Health", Nov 2014, Uppsala, Sweden. 1 p. hal-02738904

HAL Id: hal-02738904

<https://hal.inrae.fr/hal-02738904>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Title:**Toward the development of functional foods for elderly****Authors & affiliations:**

A. PROMEYRAT, T. ASTRUC, A., VENIEN, C. FERREIRA and V. SANTE-LHOUELIER


INRA, UR370 *Qualité des Produits Animaux*, 63122 Saint Genès Champanelle, France.

Extensive research deals on the development of functional foods. These foods claim to improve human health by providing benefits that going beyond the basic nutritional functions [1]. In this context, this study aims to design a food model which responds to physiological needs of elderly *i*) by a high content in easily digestible proteins *ii*) and the addition of ingredients that may reduce chronic disease risk *iii*) while providing a taste and aroma adapted to ageing of sensory function.

Several meat emulsions, Frankfurter-type, were manufactured with pork lean and backfat coming from animals feed with traditional plant sources rich in omega 3. Meat proteins are efficient to stimulate muscle protein synthesis in elderly subjects due to a favourable balance in indispensable amino acids and their high digestibility [2].

Five different meat emulsions were prepared to assess the effect of *i*) the cutting duration, *ii*) the low fat content and *iii*) the addition of psyllium or tapioca starch, on the size, the form and the distribution of fat droplets (Fig. 1). Psyllium was added for its health benefit in prevention to chronic diseases and for its emulsifying power [3]. Tapioca starch was used as a fat replacer in order to improve the flavour and juiciness by its capacity to retain moisture [4]. Changes of fat droplets, known to alter the sensory perception in mouth [5], were determined using microscopy techniques.

a)	C/LF	C/HF	F	F/FP	F/TS
Cutting duration	5 min	5 min 3	11 min	8 min 3	8 min 3
Backfat	14	27	14	14	14
Meat	66	54	66	66	66
Plasma	3	3	3	3	3
Nitrite salt	2	2	2	2	2
Spices	0.3	0.3	0.3	0.3	0.3
Lactose	0.5	0.5	0.5	0.5	0.5
Ice	14	14	14	11	11
Psyllium	0	0	0	3	0
Tapioca Starch	0	0	0	0	3



b) Fat droplet (Measurements of size and form of droplets)

C/LF, control with low fat content; C/HF, control with high fat content ; F, Frankfurters with long time of cutting ; F/FP, Frankfurters + psyllium fibers; F/TS, Frankfurters + tapioca starch.

Fig. 1: (a) Formulation of Frankfurters (g/100g); (b) Determination of the size and form of the fat droplets using light microscopy (red oil staining).

The fat droplets were strongly affected by the cutting duration. An increase of cutting time reduced significantly the size of fat droplets. In contrast, the fat droplets were slightly altered by the reduction of fat content and the addition of psyllium or tapioca starch. This research allowed developing a relevant food model in combining technological, nutritional and organoleptic considerations.

- [1] Hasler, C. M. (2002). Functional foods: Benefits, concerns and challenges-a position paper from the American Council on Science and Health. *Journal of Nutrition*, 132(12), 3772-3781.
- [2] Bax, M.-L., Buffière, C., Hafnaoui, N., Gaudichon, C., Savary-Auzeloux, I., Dardevet, D., Santé-Lhoutellier, V., & Rémond, D. (2013). Effects of Meat Cooking, and of Ingested Amount, on Protein Digestion Speed and Entry of Residual Proteins into the Colon: A Study in Minipigs. *PLoS ONE*, 8(4), e61252.
- [3] Yu, L., Lutterodt, H., & Cheng, Z. (2008). Chapter 4 Beneficial Health Properties of Psyllium and Approaches to Improve Its Functionalities. In L. T. Steve (Ed.), *Advances in Food and Nutrition Research*, vol. Volume 55 (pp. 193-220): Academic Press.
- [4] Hughes, E., Mullen, A. M., & Troy, D. J. (1998). Effects of fat level, tapioca starch and whey protein on frankfurters formulated with 5% and 12% fat. *Meat Science*, 48(162), 169-180.
- [5] van Ruth, S. M., & Roozen, J. P. (2000). Influence of mastication and saliva on aroma release in a model mouth system. *Food Chemistry*, 71(3), 339-345.