



What the brain tells us about food neophobia in the young chicken *Gallus gallus domesticus*

Emilie Perez, Marion Georgelin, Paul Constantin, Fabien Cornilleau, Maryse Meurisse, Aline Bertin, Ludovic Calandreau

► To cite this version:

Emilie Perez, Marion Georgelin, Paul Constantin, Fabien Cornilleau, Maryse Meurisse, et al.. What the brain tells us about food neophobia in the young chicken *Gallus gallus domesticus*. 8. European Conference on Behavioural Biology (ECBB), Jul 2016, Vienne, Austria. , 2016, 8th European Conference on Behavioural Biology (ECBB). hal-02739054

HAL Id: hal-02739054

<https://hal.inrae.fr/hal-02739054>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A46. What the brain tells us about food neophobia in the young chicken *Gallus gallus domesticus*

Emilie C. Perez , Marion Georgelin, Paul Constantin, Fabien Cornilleau, Maryse Meurisse, Aline Bertin and Ludovic Calandrea

Institut National de la Recherche Agronomique INRA, UMR85 Physiologie de la Reproduction et des Comportements, Nouzilly, France; CNRS, UMR6175 F-37380 Nouzilly, France; Université de Tours, F-37041 Tours, France; Haras Nationaux, F-37380 Nouzilly, France

Food neophobia is a complex behaviour that involves the complete or partial rejection to eat a new source of food. While described in a large variety of mammalian and bird species in terms of behavioural changes, there are still very few studies to date interested in its neurobiological bases. However, understanding the neuronal bases of food neophobia will bring new insights to reduce this behaviour in farm birds, thus increasing their welfare. We propose here for the first time a comprehensive study in the domestic chicken, from a detailed description of the behaviour to the neural activity associated with food neophobia. Using both behavioural and immunohistochemistry procedures on one-week chicks, our study focuses on several brain structures suspected to play an important role in food neophobia, including: five amygdala structures implicated in emotions, the nucleus accumbens that plays a central role in the reward circuit, and the olfactory bulb so as to understand the possible olfactory dimension of the behaviour.