



Multi-varietal genomic selection in French pig populations

Céline Carillier-Jacquin, Llibertat Tusell, Juliette Riquet, Marie-José Mercat,
Catherine Larzul

► To cite this version:

Céline Carillier-Jacquin, Llibertat Tusell, Juliette Riquet, Marie-José Mercat, Catherine Larzul. Multi-varietal genomic selection in French pig populations. 67. Annual Meeting of the European Federation of Animal Science (EAAP), Aug 2016, Belfast, United Kingdom. 721 p., 10.3920/978-90-8686-284-9 . hal-02739126

HAL Id: hal-02739126

<https://hal.inrae.fr/hal-02739126>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Multi-varietal genomic selection in French pig populations

Céline Carillier-Jacquin

Llibertat Tusell Palomero

Juliette Riquet

Marie-José Mercat

Catherine Larzul

INRA-GenPhySE, Toulouse, France



Context

- **Small population size = lower GEBV accuracy**
- **Nucleus from different breeding organizations > same breed but different populations**
- **Including animals from a biggest population to small one = multi-varietal genomic evaluation**
- **Objective : look if multi-varietal evaluation could improve accuracy**

Data collection



- 3 varieties of Piétrain pig genotyped:

- V2 and V3 have limited population size

- Test multi-varietal genomic evaluations: V1+V2 and V1+V3

	Sires	Offspring
V1	96	752
V2	13	118
V3	17	177

Single step genomic evaluation

- Single step genomic evaluation using blupf90 program
- Phenotype of 3 varieties considered as the same trait
- Genetic parameter used = estimated in V1
- 60 traits measured (production, blood and hormonal parameters, skin lesions)

	h^2
Growth rate	0.6
Feed conversion ratio	0.4
Androstenone	0.6
Total number of skin lesions	0.3

Estimation of prediction ability

- 4-fold cross validation
- In 7 cases :

Reference population	V1	V2	V3	V1+V2	V1+V2	V1 + V3	V1+V3
Validation set	V1 sires	V2 sires	V3 sires	V1 sires	V2 sires	V1 sires	V3 sires

- Prediction ability = $\text{corr}(\text{GEBVpred}, \text{GEBVtrue})$ for validation set
- Total prediction ability = mean of the 4 prediction abilities

Prediction ability for V2 and V3 sires


Prediction ability

0.70
0.60
0.50
0.40
0.30
0.20
0.10
0.00

Prediction ability


0.70
0.60
0.50
0.40
0.30
0.20
0.10
0.00

Prediction ability for V1sires




Sires V1
with the largest reference population size

Estimation of theoretical accuracy



$$\text{accuracy(V2sires)} = \sqrt{\frac{\sigma_u^2 - IPEW}{\sigma_u^2}}$$

Estimation of theoretical accuracy





$$\text{accuracy}(\text{siresV1}) = \sqrt{\frac{\sigma_u^2 - PEV}{\sigma_u^2}}$$

$$\text{accuracy}(\text{siresV3}) = \sqrt{\frac{\sigma_u^2 - PEV}{\sigma_u^2}}$$

.09

Similar results for theoretical accuracies of sires



.010

Multi-varietal genomic evaluation could improve accuracies

- Improvement in prediction ability:
 - from +1.6% to +322% for V3 sires
 - from +1.2% to +261% for V2 sires

kinship coeff(V1 and V3) = 12%
kinship coeff(V1 and V2) = 8%

- enhancement proportional to the degree of relatedness
- No improvement/degradation for V1 sires

 Multi-varietal evaluation improve accuracy for small populations

Thank you for your attention!

- Utopige fundings



- Bioporc (ADN, Choice Genetics France, Gene +, Nucleus)



- IFIP and Le Rheu test station staff



- Ignacy Misztal for blupf90 suite of programs



Traits recorded

	h^2
Back fat thickness	0.6
Muscle thickness	0.3
Daily feed intake	0.4
Ph in semimembranosus	0.2
Ph in LD*	0.2
Drip loss	0.1
Dressing yield	0.5
Total	0.6
On right part	0.03
On left part	0.02
Number of lesion on carcass	0.12
On front part	0.3
On rear part	


	h^2
Red indices in GM*	0.5
Red indices in GS*	0.6
Red indices in LD*	0.7
Yellow indices in GM*	0.6
Yellow indices in GS*	0.1
Yellow indices in LD*	0.5
Lightness in GM*	0.1
Lightness in GS*	0.7
Lightness in LD*	0.2
Back fat weight	0.8
Percent of ham cut	0.5
Belly weight	0.3
Percent of loin	0.5
Percent of shoulder	0.3

Traits recorded


	h^2
Blood volume	0.9
C-Reactive protein	0.3
Pig map*	0.2
Estradiol level	0.3
Hematocrit	0.4
Blood count	0.2
Indole level	0.7
Number of leucocytes	0.2
Intramuscular fat	0.4
number of lymphocytes	0.2
Number of pellets	0.3
Skatol level	0.2
Testosterone level	0.3
Volume of pellets	0.6

	h^2
Number of lesion at the beginning of growing	Total
	Total on one side
	Total on the other side
	Total on front part
	Total on rear part
Number of lesion at the end of growing	Total
	Total on one side
	Total on the other side
	Total on front part
	Total on rear part

4-fold cross validation




4-fold cross validation




Total prediction ability (sire V1) = mean of the 4 prediction abilities

4-fold cross validation



4-fold cross validation



Total prediction ability (sire V1) = mean of the 4 prediction abilities

4-fold cross validation

