



HAL
open science

Modelling anaerobic digestion and compost of organic residues: towards organic matter fate prediction in soils

Julie Jimenez, Quentin Aemig, Jean-Philippe Steyer, Dominique Patureau,
Sabine Houot

► To cite this version:

Julie Jimenez, Quentin Aemig, Jean-Philippe Steyer, Dominique Patureau, Sabine Houot. Modelling anaerobic digestion and compost of organic residues: towards organic matter fate prediction in soils. ORBIT 2016 Organic Resources and Biological Treatment - 10. International Conference on Circular Economy and Organic Waste, Technological and Educational Institute of Crete (TEI of Crete). GRC., May 2016, Heraklion, Greece. 200 p. hal-02739210

HAL Id: hal-02739210

<https://hal.inrae.fr/hal-02739210>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

MODELLING ANAEROBIC DIGESTION AND COMPOST OF ORGANIC RESIDUES: TOWARDS ORGANIC MATTER FATE PREDICTION IN SOILS

J. Jimenez^{1*}, Q. Aemig¹, J-P. Steyer¹, S. Houot², D. Patureau¹

¹INRA, UR0050, Laboratoire de Biotechnologie de l'Environnement, Narbonne, F-11100, France, ²INRA UMR INRA-AgroParisTech Ecologie et Ecotoxicologie des Agroecosystemes, 78 850 Thiverval-Grignon, France

Contact: Dr. Julie Jimenez, INRA LBE, avenue des Etangs 11100 Narbonne France, +33 4 68 42 51 83, julie.jimenez@supagro.inra.fr

EXECUTIVE SUMMARY

In a context of environmental biorefinery, organic residues such as sludge, biowaste, manure are no more considered as wastes but as potential resources for some targeted services. These services encompass the energy production through the methane produced by their anaerobic digestion, the cropped soils fertilisers and amendments production from digestates, and other value-added compounds production (e.g. organic acids and hydrogen obtained during fermentation). There are currently only few studies focused on combined optimization of energetic and agronomic values of organic wastes. One way to optimize both objectives is through process modelling. However, input variables from all existing models are different and there is a need to find relevant and shared input variables to model the fate of organic residues during their treatments. Recently, a new promising methodology of organic matter characterization has been successfully used to predict anaerobic and aerobic biodegradability and bioaccessibility of a broad range of organic residues (Jimenez *et al.*, 2015). This methodology is based on the combination of chemical fractionation simulating organic matter accessibility and of a 3D fluorescence spectroscopy highlighting fraction's complexity. The fractions obtained were used as models input variables. Existing models (Batstone *et al.*, 2002, Zhang *et al.*, 2012, Garnier *et al.*, 2003) have been modified to take into account these new variables. In order to calibrate and validate the models, a lab scale reactor of anaerobic digestion was operated to treat wastewater sludge. Biogas, methane proportion, pH and input and output digestate quality were monitored. After 60 days of steady state operation, the digestate was then centrifuged to separate the liquid from the solid part. The solid part was then added with green waste into a lab scale compost reactor. Temperature, CO₂ production, moisture and compost quality were monitored. Finally, the final compost was used as substrate for cropped soil incubation. First results showed that, the new anaerobic digestion model was successfully able to predict biogas and digestate quality when model and experimental data were compared. The final quality of the solid part of the simulated digestate was then introduced with the green waste fractionation to model composting. The modified compost model also fitted the quality of compost during the different steps of the composting. Concerning soil incubation modelling, experimental results are still ongoing but will be part of the oral presentation. Overall, the obtained results constitute a strong basis for the development of a decision support system to optimize the design and operation of organic wastes treatment plants.

REFERENCES

- Batstone, D. J., Keller, J., Angelidaki, I., Kalyuzhnyi, S. V., Pavlostathis, S. G., Rozzi, A., Sanders, W. T. M., Siegrist, H., Vavilin, V. A. (2002) *Anaerobic Digestion Model No.1. (ADM1). IWA Scientific and Technical Report No. 13*. IWA, ISBN:1-900222-78-7.
- Garnier, P., Neel C., Aita, C., Recous, S., Lafolie, F., Mary, B., (2003). *Modelling carbon and nitrogen dynamics in a bare soil with and without straw incorporation*. European journal of soil science 54, pp. 555-568.
- Jimenez, J., Aemig, Q., Doussiet, N., Feurgard, I., Steyer, J.-P., Patureau, D., Houot, S. (2015): *Towards a unified organic waste characterization of organic matter accessibility and complexity*. Bioresource Technology 194, pp.344-353.
- Zhang, Y., Lashermes, G., Houot, S., Doublet, J., Steyer, J.P., Zhu, Y.G., Barriuso, E., Garnier, P., (2012). *Modelling of organic matter dynamics during the composting process*. Waste Manag. 32, 19–30.