



HAL
open science

Do competition and predation risk affect weed seed selectivity in granivorous carabid beetles ?

Alice Charalabidis, David Bohan, Sandrine Petit, F.X.

Dechaume-Moncharmont

► To cite this version:

Alice Charalabidis, David Bohan, Sandrine Petit, F.X. Dechaume-Moncharmont. Do competition and predation risk affect weed seed selectivity in granivorous carabid beetles ?. International Conference of Ecological Sciences Sféologie-2016, Société France d'Ecologie (FFSPN). FRA., Oct 2016, Marseille, France. hal-02739560

HAL Id: hal-02739560

<https://hal.inrae.fr/hal-02739560>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Do competition and predation risk affect weed seed selectivity in granivorous carabid beetles?

Alice Charalabidis^{*1}, David Bohan¹, Sandrine Petit¹, and François-Xavier Dechaume-Moncharmont²

¹Agroécologie – Institut national de la recherche agronomique (INRA) : UMR1347, Université de Bourgogne – 17 rue Sully, BP 86510, F-21000 Dijon, France

²Biogéosciences – Université de Bourgogne, CNRS : UMR6282 – 6 Bvd Gabriel 21000 DIJON, France

Abstract

Carabid beetles consume substantial amounts of weed seeds and appear to be good candidates for the biological control of weeds in arable fields. Yet, their selection criteria for certain seeds are still not clearly understood.

When foraging in-field, individuals have to deal with competition and predation exerted by conspecifics or other species of carabid beetles; they, thus, often face a trade-off between staying safe (i.e. avoiding predation or starvation) and taking the time to assess the quality of the food. We hypothesize that, in order to maximise their fitness, carabid individuals will modify their pattern of food selection (choosiness) while foraging under varying levels of risk.

To test this hypothesis, the foraging behaviour of the seed feeding carabid beetle *Harpalus affinis* was examined under 4 different experimental conditions: i) predation risk; ii) intraspecific competition; iii) interspecific competition; and, iv) control. All the risks were simulated using chemical cues from conspecific beetles or different species of carabid beetles (predatory or granivorous).

Our results show that when foraging under a risk of predation or competition for food, *H. affinis* individuals reduced their choosiness, and that the risk of predation affected choosiness twice as much as competition. This finding suggests that *H. affinis* individuals are able to finely adjust their choosiness in response to the level of risk they face, potentially in order to maintain their food intake while reducing their exposure to risk

Keywords: predator, prey, intraspecific competition, interspecific competition, choosiness, carabid beetles

^{*}Speaker