



HAL
open science

Combining genetic resistance and management of field margins to control virus epidemics in melon crops

Alexandra Schoeny, Cecile Desbiez, Catherine Wipf-Scheibel, Pauline Millot,
Karine Nozeran, Patrick Gognalons, Hervé Lecoq

► To cite this version:

Alexandra Schoeny, Cecile Desbiez, Catherine Wipf-Scheibel, Pauline Millot, Karine Nozeran, et al.. Combining genetic resistance and management of field margins to control virus epidemics in melon crops. 13. International plant virus epidemiology symposium, Jun 2016, Avignon, France. 165 p. hal-02739628

HAL Id: hal-02739628

<https://hal.inrae.fr/hal-02739628v1>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

COMBINING GENETIC RESISTANCE AND MANAGEMENT OF FIELD MARGINS TO CONTROL VIRUS EPIDEMICS IN MELON CROPS

Schoeny, A., Desbiez, C., Wipf-Scheibel, C., Millot, P., Nozeran, K., Gognalons, P., and Lecoq, H.

INRA, UR407 Pathologie Végétale, CS 60064, 84143 Montfavet Cedex, France
(alexandra.schoeny@avignon.inra.fr)



BACKGROUND and OBJECTIVES

The melon *Vat* gene confers resistance to the aphid species *Aphis gossypii* and resistance to non-persistent viruses (CMV, WMV...) when inoculated by this species (1). It is however ineffective in blocking the transmission of viruses inoculated by other aphid species. Its effect on non-persistent virus epidemics is thus expected to be limited when *A. gossypii* is not the main aphid species visiting the melon crop. On the contrary, although not documented, its effect on the persistent virus CABYV mainly vectored by *A. gossypii* is expected to be strong.

The first objective of the study was to evaluate the effect of *Vat* on virus epidemics and on the genetic structure of virus populations in field conditions.

The second objective was to investigate the benefit of combining *Vat* resistance and an appropriate management of field margins to regulate the populations of aphids and/or their virus load. Indeed, literature suggests that flower strips can participate in pest biological control by favoring natural enemies (2), and strips of non-host plants can protect crops from non-persistent viruses by allowing aphids to probe on healthy plants and thus to lose their virus load before reaching the crops (3).

MATERIALS and METHODS

A five-year field experiment was conducted to compare two types of field margin management (bare soil and flower strip) on the efficiency of *Vat*. Virus epidemics were monitored by DAS-ELISA and genetic structures of virus populations characterized by RT-PCR with specific primers and sequencing. The effect of *Vat* and field margin management on the infection probability of a plant by a given virus was calculated with a generalized linear model (binomial distribution and logit link function).

RESULTS

Vat reduced the development of CABYV epidemics significantly every year. Concerning non-persistent viruses, it had no effect on WMV epidemics, but allowed to reduce CMV epidemics some years, confirming the role of *A. gossypii* in CMV transmission. The genetic structure of virus populations was not affected by *Vat*. The presence of flower strips had a dual effect on virus epidemics: mostly beneficial for WMV but sometimes detrimental for CMV.

CONCLUSIONS

Combining genetic resistance and management of field margins appeared generally as a promising way to decrease the risk of viral epidemics. Nevertheless, the presence of natural enemies enhanced by flower strips may also have a detrimental effect on non-persistent virus dispersion through a modification of aphid behaviour.

REFERENCES

- (1) Boissot N, Thomas S, Chovelon V and Lecoq H (2016). NBS-LRR-mediated resistance triggered by aphids: viruses do not adapt; aphids adapt via different mechanisms. *BMC Plant Biology* 16:25
- (2) Pfiffner L and Wyss E (2004). Use of sown wildflower strips to enhance natural enemies of agricultural pests. In: G M Gurr, S D Wratten & M A Altieri (eds.), *Ecological engineering for pest management. Advances in habitat manipulation for arthropods*. CSIRO Publishing. pp. 167-188
- (3) Hooks CRE and Fereres A (2006). Protecting crops from non-persistently aphid-transmitted viruses: a review on the use of barrier plants as a management tool. *Virus Research*, 120:1-16

Building bridges between disciplines for sustainable management of plant virus diseases



13th International Plant Virus Epidemiology Symposium
6-10 June 2016, Avignon, FRANCE

Programme and Abstracts