



HAL
open science

Joint estimation of effective population size and selection coefficient without neutral markers: method validation and application to experimental evolution of viruses

Elsa Rousseau, Benoît Moury, Ludovic Mailleret, Rachid R. Senoussi, Alain Palloix, Vincent Simon, Frédéric Grognard, Sophie Valière, Frédéric Fabre

► To cite this version:

Elsa Rousseau, Benoît Moury, Ludovic Mailleret, Rachid R. Senoussi, Alain Palloix, et al.. Joint estimation of effective population size and selection coefficient without neutral markers: method validation and application to experimental evolution of viruses. 13. International plant virus epidemiology symposium, Jun 2016, Avignon, France. 165 p. hal-02739698

HAL Id: hal-02739698


<https://hal.inrae.fr/hal-02739698v1>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Building bridges between disciplines for sustainable management of plant virus diseases



IPVE 2016

13th International Plant Virus Epidemiology Symposium
6-10 June 2016, Avignon, FRANCE

Programme and Abstracts

JOINT ESTIMATION OF EFFECTIVE POPULATION SIZE AND SELECTION COEFFICIENT WITHOUT NEUTRAL MARKERS: METHOD VALIDATION AND APPLICATION TO EXPERIMENTAL EVOLUTION OF VIRUSES

Elsa Rousseau (1,2,3), Benoît Moury (3), Ludovic Mailleret (1,2), Rachid Senoussi (4), Alain Palloix (5), Vincent Simon (3,6), Frédéric Grognard (1), Sophie Valière (8) and Frédéric Fabre (9)

(1) Biocore Team, INRIA, Sophia Antipolis, France

(2) UMR ISA, CNRS, INRA, Université Nice Sophia Antipolis, Sophia Antipolis, France

(3) UR PV, INRA, Montfavet, France

(4) UR BioSp, INRA, Avignon, France

(5) UR GAFL, INRA, Montfavet, France

(6) UMR BFP, INRA, Villenave d'Ornon, France

(7) GeT-PlaGe, INRA, Genotoul, Castanet-tolosan, France

(8) UAR DEPT GA, INRA, Castanet-Tolosan, France

(9) UMR SAVE, INRA, Villenave d'Ornon, France (frederic.fabre@bordeaux.inra.fr)

BACKGROUND and OBJECTIVES

Experimental evolution studies deserve considerable attention to the estimation of basic evolutionary forces such as selection and genetic drift. With the advent of high-throughput sequencing techniques, these studies gained a renewed attention. However the joint estimation of selection and genetic drift still remain challenging when no neutral markers are available, a common situation with many microbes, such as viruses, due to their small tightly packed genomes.

MATERIAL and METHODS

We have developed a mechanistic-statistical model for estimating effective population size N_e and selection coefficient s from genetic time series data generated in evolve-and-resequence experiments. Numerical simulations of the dynamics of competing genotypes in asexual haploid Wright-Fisher populations subjected to contrasting genetic drift ($N_e \in [20, 5000]$) and selection ($|s| \in [0, 0.1]$) regimes were used to test the model. This model was then applied to data for competition between five variants of *Potato virus Y* (PVY, genus *Potyvirus*) in 15 doubled haploid lines of pepper (*Capsicum annuum*, family *Solanaceae*) that constituted different host environments. The frequencies of the variants were determined at six time-points, in eight plants per time point for each host environment.

RESULTS

The numerical experiments validated the method and made it possible to obtain bias-adjusted estimators of s for each competing genotype and of the dynamics of N_e over the time course of the experiment ($R^2 \in [0.93, 0.95]$, slopes $\in [0.98, 1.07]$ and intercepts $\in [-0.14, 0.06]$ for the best linear fit between true and estimated parameters). The real-life experiment showed that the virus populations experienced either stochastic or deterministic evolution, depending on host genotype and that N_e and s were genetically controlled by the host genotype.

CONCLUSIONS

This method constitutes an advance in the joint estimation of key population genetic parameters from time-sampled data in conditions in which no neutral markers are available and both selection and genetic drift are strong. This is a situation frequently encountered in studies of microorganisms. The observation of highly contrasted genetic drift between plant genotypes may have practical implications for slowing viral emergence through the careful choice of plant cultivars.