



HAL
open science

Variation and origin of the tomato

J. Blanca, J. Montero-Pau, Christopher Sauvage, Guillaume Bauchet, E. Illa,
M. Diez, D. Francis, Mathilde M. Causse, E. van Der Knaap, Joaquin
Canizares

► **To cite this version:**

J. Blanca, J. Montero-Pau, Christopher Sauvage, Guillaume Bauchet, E. Illa, et al.. Variation and origin of the tomato. Applied Vegetable Genomics, Vienna Institute for Nature Conservation and Analyses (VINCA). AUT., Feb 2014, Vienne, Austria. pp.12. hal-02739830


HAL Id: hal-02739830

<https://hal.inrae.fr/hal-02739830>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



VISCEA

Vienna International Science
Conferences and Events Association




International Conference

Applied Vegetable Genomics

Programme and Abstracts

Vienna, Austria

19 - 20 February 2014



Variation and Origin of the Tomato

**J. Blanca¹, J. Montero-Pau¹, C. Sauvage², G. Bauchet², E. Illa³, M. Díez¹, D. Francis³, M. Causse²,
E. van der Knaap³, J. Cañizares¹**

¹COMAV, Polytechnic University of Valencia, Camino de Vera s/n, 46022, Valencia, Spain, ²GAFI-INRA, Domaine Saint-Maurice, F-84143 Montfavet Cedex, France, ³Department of Horticulture and Crop Science, Ohio Agricultural Research and Development Center (OSU), Wooster, OH 44691, USA

A joint SolCAP dataset of 1000 tomato genotypes was assembled including a comprehensive representation of *S. pimpinellifolium* (SP), *S. lycopersicum* varieties cerasiforme (SLC) and *lycopersicum* (SLL). Traditionally SLC has been considered as formed by the cherry tomatoes, but that was shown to be a misconception. The cherry tomatoes included both SLC accessions and hybrids and admixtures between SP, SLC and SLL. The results showed that SLC is a semi-domesticated with a key role in the complex history of the tomato domestication. Moreover SLC contained the South American vintage cultivated tomatoes. Andean SLC had a gene diversity comparable to SP, while non-Andean SLC and vintage SLL showed a lower diversity. The contemporary cultivated SLL also had a low diversity despite the introgressions added in the second half of the 20th century.