



Riboflavin secretion in cow's milk varies according to diet composition and season

Sophie Laverroux, Julie Vallet, Chantal Chassaing, Christiane Girard, Claire Agabriel, Bruno Martin, Benoit Graulet

► To cite this version:

Sophie Laverroux, Julie Vallet, Chantal Chassaing, Christiane Girard, Claire Agabriel, et al.. Riboflavin secretion in cow's milk varies according to diet composition and season. 1. Joint Meeting of FAO-CIHEAM Mountain Pastures and Mediterranean Forages Resources Networks and Mountain Cheese Network, Jun 2014, Lempdes, France. pp.323-326. hal-02739841

HAL Id: hal-02739841

<https://hal.inrae.fr/hal-02739841>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Riboflavin secretion in cow's milk varies according to diet composition and season

S. Laverroux^{1,2}, J. Vallet^{1,2}, C. Chassaing^{2,1}, C.L. Girard³, C. Agabriel^{2,1}, B. Martin^{1,2}, B. Graulet^{1,2}

¹ Inra, UMR1213 Herbivores, F-63122, Saint-Genès-Champagnelle (France)

² Clermont Université, VetAgro Sup, UMR1213 Herbivores, BP 10448, F-63000, Clermont-Ferrand (France)

³ Centre de Recherche sur le Bovin Laitier et le Porc, Agriculture et Agroalimentaire Canada, Sherbrooke, (Canada)

Abstract. In ruminants, B-vitamin supply from the diet and synthesis by the rumen microflora is considered sufficient to avoid deficiency. Nevertheless, factors affecting their concentrations in milk have never been studied in-depth, even if dairy products are among the best contributors of several vitamins (especially riboflavin) in human nutrition. We compared, during a year, riboflavin concentrations in bulk tank milks from 20 groups of 5 commercial farms divided among production systems mainly characterized by the forage nature (grass or maize silage). Riboflavin concentrations were greater in milks from grass-based than maize silage systems (1.91 vs 1.70 mg.L⁻¹, P<0.001). Moreover, they increased during summer (from 1.71 to 1.99 mg.L⁻¹, P<0.001) whatever the feeding system. These differences were largely explained by a dilution effect due to the level of the milk yield but the mean amount of riboflavin secreted in milk per cow per day was higher when cows were fed a diet rich in maize silage by comparison to a herb-based diet (42.5 vs 33.4 µg.d⁻¹, respectively, P< 0.001). Conversely to what should be thought, riboflavin concentrations in milk are submitted to variations, and more interestingly regulations, that influence the global nutritional quality of milk.

Keywords. Cow milk – vitamin – feeding system - season.

Title. *La sécrétion de riboflavine dans le lait de vache varie selon la composition de la ration et la saison*

Résumé. En raison de leur double origine alimentaire et microbienne ruminale, la disponibilité en vitamines B chez les ruminants est considérée pouvoir couvrir les besoins de ces animaux. C'est la raison pour laquelle, la variabilité de leurs concentrations dans le lait n'a jamais été explorée, même si le lait et les produits laitiers sont parmi les premiers contributeurs de certaines vitamines (dont la riboflavine) en nutrition humaine. Nous avons déterminé les concentrations en riboflavine dans des laits de tournées simulées correspondant à 20 groupes de 5 fermes commerciales caractérisés par leur système fourrager (basé sur l'herbe ou l'ensilage de maïs). Les concentrations de riboflavine étaient supérieures dans les laits issus des systèmes à base d'herbe par rapport à ceux utilisant préférentiellement le maïs (1.91 vs 1.70 mg.L⁻¹, P<0.001). Elles ont augmenté également pendant l'été quel que soit le système considéré (de 1.71 à 1.99 mg.L⁻¹, P<0.001). Ces différences s'expliquent principalement par un effet de dilution lié à la quantité de lait produite ; cependant, la quantité moyenne de riboflavine sécrétée dans le lait par vache et par jour était supérieure lorsque les vaches recevaient une ration à base d'ensilage de maïs que lorsqu'elles recevaient une ration à base d'herbe (respectivement 42.5 vs 33.4 µg.j⁻¹, P<0.001). Contrairement au concept couramment admis, la concentration de riboflavine du lait est variable, et de manière encore plus intéressante, elle semble être régulée, ce qui impacte la qualité nutritionnelle globale du lait et des produits laitiers.

Mots-clés. Lait de vache – vitamine – système d'alimentation - saison.

I – Introduction

For decades, availability of B vitamins has been globally considered as sufficient to cover the estimated requirements of dairy cows (except vitamins B₅ and B₉) thanks to supply from the diet and/or from synthesis by rumen microorganisms (National Research Council, 2001). However, as milk production levels considerably increased through bovine genetic selection and optimised feed efficiency, it is obvious that cow requirements for B vitamins, which are involved

in the main metabolic pathways (cell respiration, energy production, nucleic acid synthesis, amino-acid metabolism and protein synthesis, neoglucogenesis...), increased proportionally (Girard et al., 2010). To date, the relationship between performance levels and B vitamins requirements has not been studied. Proofs that B vitamin availability to lactating cows would be lower than needs are insufficient even though some experimental results suggested it. For example, increasing B vitamin supply (especially B₉ and B₁₂) to lactating dairy cows has beneficial effects, including on milk performance (Girard et al., 2010). Moreover, it was also observed that their dietary supplementation raised their concentration in milk (Graulet et al., 2007). This latter observation was interesting from a nutritional point of view, because milk and dairy products are good sources of vitamin A, B₁₂ (cobalamins) and B₂ (riboflavin) (Coudray, 2011). However, the factors regulating concentration of B vitamins in milk have not been fully characterized. In a first publication, we reported that the highest levels of milk vitamin B₉ and vitamin B₁₂ were generally associated with grass- and maize silage-based diets, respectively (Chassaing et al., 2011). In the present work, we compared riboflavin secretion in milks obtained from different production conditions varying in term of diet composition, season or altitude.

II – Materials and methods


The study was carried out in 20 groups of five farms divided among four production systems mainly characterized by their forage system and altitude: feeding systems based on grass in Mountain (GM) or in Lowland (GL) and maize silage feeding systems located in Mountain (MM) or in Lowland (ML). In GM and GL, forages were mainly hay during the wintering period and fresh grass during the grazing period. The proportion of maize in cow diets was lower in MM than in ML whatever the period of the year. It was particularly marked in summer when the proportion of maize silage was extremely reduced in MM at the benefit of pasture / grass silage or grass silage / hay combination. In each group of farms, bulk tank milk was sampled five times in the course of the year 2008 at key times in animal feeding patterns: twice in the overwintering period with diets based on preserved forage (January and February), and three times during the grazing period: in May, July and September. During the week of each milk sampling, a survey was carried out to record herd characteristics, performance and feeding. Average diets for each group were described on the basis of proportions of forages and concentrate in the diet calculated from the declared quantities the farmers dispensed. When the amount of one feed ingested was unknown (such as pasture), the estimation of the amount ingested was based on the energy requirements. The energy balance was supposed equal to zero. Milk riboflavin was extracted in duplicate by successive acid hydrolysis and enzymatic treatments using papain and acid phosphatase. Vitamin concentrations were measured in the extracts using a Acquity UPLC equipped with a 150 x 2.1 mm HSS T3, 1.8- μ m column (Waters, France) and fluorescence detection (λ_{exc} = 400 nm ; λ_{em} = 520 nm). Data were processed by Anova using the MiniTab16 software introducing the production system, period, and the interaction into the model as fixed effects. Because the characteristics of the production systems changed between periods, especially the diet composition, period was not considered a repeated factor.

III – Results and discussion

Riboflavin concentrations in the hundred milks studied varied from 1.51 to 2.15 mg.L⁻¹ with a mean value of 1.78 ± 0.02 mg.L⁻¹. As for vitamin B₉ (Chassaing et al., 2011), riboflavin concentrations were greater in milks from grass-based than maize silage systems (1.91 vs 1.70 mg.L⁻¹, P<0.001, Fig. 1A). However, the interaction between diet composition and altitude was significant (P<0.01) because riboflavin concentrations were the highest in milks produced in lowland grass-based systems whereas the lowest values were observed in maize silage based systems also in lowland. Whatever the production systems, riboflavin concentrations increased

during summer (from 1.71 to 1.99 mg.L⁻¹, P<0.001). Values observed in maize silage systems were in agreement with food tables (Graulet et al., 2013). However, riboflavin concentrations in milks produced in a grass-based system were higher (+14%), reinforcing the nutritional value of these milks.

Figure 1: Variations in milk riboflavin concentration (A) and secretion (B) according to feeding system, altitude and season



Production System : GM: Grass Mountain (---△---) ; GL: Grass Lowland (—▲—) ;
MM: Maize Mountain (---□---) ; ML: Maize Lowland (—■—).
Means with different superscripts differ (P<0.05).

Riboflavin secretion in milk per day and per cow was estimated for each system using the corresponding mean milk yield. Milk production levels were largely higher in maize-silage than in grass-based systems, especially in winter and in lowland. Consequently, the amount of riboflavin secreted in milk per day per cow was higher in maize- than in grass-based systems (42.5 vs 33.4 $\mu\text{g}\cdot\text{d}^{-1}$, respectively, $P < 0.001$, Fig.1B). However, it was almost stable during the year and equivalent between lowland and mountain systems. Thus, variations in riboflavin concentrations in milk were largely explained by a dilution effect due to the higher milk yield in maize based-systems. The effects of season and altitude on the total amount of riboflavin secreted in milk are negligible. However these data suggest that riboflavin supply is greater for cows fed maize silage than for cows fed grass-rich diets, as illustrated by the amounts of vitamin secreted by the mammary gland. Of course, riboflavin could be provided to cows directly by the diet or indirectly by the influence of the diet on rumen fermentations, that is to be studied through complementary experiments. Finally, this last observation raise a concern about the adequacy of riboflavin supply to cows fed grass-rich diets and its role as a limiting factor for production efficiency.

IV – Conclusions

These results complete those obtained on folates (B_9) and vitamin B_{12} (Chassaing et al., 2011) confirming that milk concentrations of B vitamins vary according to diet and seasons even though complementary studies are needed for the other B vitamins and to better understand the underlying mechanisms. More interesting is the fact that riboflavin concentrations in milks produced by cows fed grass-rich diets especially in summer are higher than reference values reported in tables, reinforcing the interest of milk as a riboflavin source in human nutrition in general and giving an added value to milks produced in grass -based systems.

Acknowledgments

This work was permitted by an experimental design included in the TRUEFOOD project co-funded by the European Commission within the 6th Framework Programme. We acknowledge C. Sibra (VetAgro Sup), I. Constant (INRA), C. Fargeix and all our partners of the dairy sector and the farmers involved in this study.

References

- Chassaing C., Graulet B., Agabriel C., Martin B., Girard C.L., 2011. Vitamin B_9 and B_{12} Contents in Cow Milk according to Production System, p35-36. *In: Proceedings of the 10th International Meeting on Mountain cheese*, 14-15 september 2011, Dronero (CN), Italy. Edited by University of Turin, Italy, ISBN 978-88-902754-5-6
- Coudray B., 2011. The contribution of dairy products to micronutrient intakes in France. *In: J. Amer. Coll. Nutr.*, 30. pp. 410S-4S.
- Girard C.L., Preynat A., Graulet B., Lapierre H., 2010. Could minor nutrients such as B vitamins alter major metabolic pathways in lactating dairy cow ? Energy and protein metabolism and nutrition. Wageningen Academic Publisher, Wageningen (NLD) ; EAAP Scientific Series 3rd International Symposium on Energy and Protein Nutrition and Metabolism (ISEP), 2010, Parma (Italie), 6-10 septembre 2010. 127, 235-243.
- Graulet B., Matte J.J., Desrochers A., Doeppel L., Palin M.F., Girard C.L. 2007. Effects of dietary supplements of folic acid and vitamin B12 on metabolism of dairy cows in early lactation. *In: J. Dairy Sci.* 90. pp.3442–3455.
- Graulet B, Martin B, Agabriel C, Girard C, 2013. Vitamins in milk. *In : Milk and dairy products in human nutrition : Production, composition and health*. Y.W. Park and G.F.W. Haenlein (Editors). (p. 200-219).
- National Research Council, 2001. Nutrient requirements of dairy cattle. 7th revised edition. National Academy Press, Washington, DC.