



HAL
open science

Rethinking plant's stomatal behavior

Nicolas Martin-StPaul, Sylvain S. Delzon, Hervé H. Cochard

► **To cite this version:**

Nicolas Martin-StPaul, Sylvain S. Delzon, Hervé H. Cochard. Rethinking plant's stomatal behavior. Xylem International Meeting, Sep 2015, Bordeaux, France. 86 p. hal-02739900

HAL Id: hal-02739900

<https://hal.inrae.fr/hal-02739900>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Rethinking plant's stomatal behavior

Martin-StPaul NK^{1*}, Delzon S² & Cochard H³.

¹ INRA, UR629 Ecologie des Forêts Méditerranéennes (URFM), Avignon, France.

² INRA, University of Bordeaux, UMR BIOGECO, F-33450 Talence, France.

³ INRA, Clermont University, UMR547 PIAF, F-63100 Clermont-Ferrand, France.

*Author for correspondence: nicolas.martin@paca.inra.fr

Abstract:

It's commonly accepted that the universal trade-off between water saving and CO₂ capture driven by stomatal conductance includes two strategies of hydraulic functioning that determine plant vulnerability to drought. Accordingly, isohydric species, that close stomata rapidly to regulate water loss, can be subjected to carbon starvation. By contrast, anisohydric species can maintain significant rates of gas exchange but are predisposed to hydraulic failure because they operate with narrower hydraulic safety margins during drought. *Despite a large research effort very few convincing evidence has been reported so far.* In this study we developed an ecophysiological model that couple stomatal and hydraulic behaviors to study the response to water deficit. Based on initial soil features, plant transpiration rate, conductance and leaf area, resistance to cavitation and osmotic potential at full turgor; the model computes soil and plant water potential, leaf turgor pressure and plant and soil conductance. Stomatal regulation is a function of leaf turgor potential. The model allows computing the dynamic of dehydration, the number of days with stomata open (*i.e.* length of the photosynthetically active period) and the plant death by hydraulic failure. The model was tested for a wide range of plant and soil type. Three major outcomes have been addressed: first, any species can behave as isohydric or anisohydric according to environmental conditions. Second, for a given safety margins, species more resistant to cavitation are surprisingly more prone to drought-induced hydraulic failure. Third, to maximize carbon capture while avoiding hydraulic failure; stomatal closure should occur within a narrow range of water potential (before ~ -4 MPa) whatever the resistance to cavitation of the species. The predictions of the model for an optimal behavior (in terms of avoiding cavitation and maximizing openness of stomata) are congruent with global databases of stomatal closure, osmotic potential and resistance to cavitation. Altogether these results reject the isohydric versus anisohydric behavior of plant stomata and their implication in drought-induced plant mortality. The model can serve as a basis to improve vegetation models of biogeochemical cycles or plant distribution models in a context of anticipating risk related to increasing drought under climate change.