



**HAL**  
open science

## Role of gas accumulation in dry anaerobic digestion of wheat straws

Elisabeth Cazier, Eric Trably, Jean-Philippe Steyer, Renaud R. Escudie

### ► To cite this version:

Elisabeth Cazier, Eric Trably, Jean-Philippe Steyer, Renaud R. Escudie. Role of gas accumulation in dry anaerobic digestion of wheat straws. 14. World Congress on Anaerobic Digestion (AD14), International Water Association (IWA). INT., Nov 2015, Viña del Mar, Chile. hal-02740021

**HAL Id: hal-02740021**

**<https://hal.inrae.fr/hal-02740021v1>**

Submitted on 2 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Role of gas accumulation in dry anaerobic digestion of wheat straws

E.A. Cazier\*, E. Trably\*\*, J.P. Steyer\*, R. Escudé\*

\*INRA, UR0050, Laboratoire de Biotechnologie de l'Environnement, Avenue des Etangs, 11100 Narbonne, France

+Corresponding author: eric.trably@supagro.inra.fr; Phone: (+33) (0) 468 425 172; Fax: (+33) (0) 468 425 160

## Abstract

In dry anaerobic digestion, biogas production is inhibited at high total solids contents. Such inhibition can likely be due to a slow diffusion of dissolved molecules into the digestate with subsequent local accumulation of inhibitory products.

In this study, we investigated the impact of H<sub>2</sub> and CO<sub>2</sub> partial pressure on dry AD. Specific batch tests operated were carried out with thin layer to limit gas diffusion. Results showed that initial H<sub>2</sub> partial pressure higher than 800 mbars had a strong inhibitory effect on methane production. The total substrate degradation yields were also impacted, indicating a direct inhibitory effect on hydrolysis. Interestingly, when H<sub>2</sub> and CO<sub>2</sub> were both added, they were consumed by homoacetogenesis/methanogenesis until CO<sub>2</sub> was completely exhausted. CO<sub>2</sub> was found to be the limiting element in case of H<sub>2</sub> inhibition. The reversibility of H<sub>2</sub> inhibition was also investigated by adding CO<sub>2</sub> at different reaction times. Results showed that hydrolysis and methanogenesis inhibitions were fully reversible after CO<sub>2</sub> addition even after 18 days of operation.

Our results give new insights on probable effect of local gas accumulation in dry AD processes.

## Keywords

Biogas; Lignocellulosic residues; High solids; Dry anaerobic digestion; Gas transfer

## INTRODUCTION

Over the past ten years, Dry Anaerobic Digestion (D-AD) has gained a wide interest in Europe especially when using agricultural waste or the organic fraction of municipal waste. In AD process, the organic matter contained in the waste is converted by anaerobic bacteria into a biogas composed of CH<sub>4</sub> (50-70%) and CO<sub>2</sub> (30-50%). The remaining organic matter is called “digestate” and can be further reused on land as fertilizer after considering sanitary and environmental impacts.

The key parameter that drives microbial processes in D-AD (>15% of total solids TS) is the low water content since microbial end-products could locally accumulate and inhibit methanogenesis (Abbassi-Guendouz et al., 2012).

Water availability depends on two forms: (1) the low available “bound water” where the water is chemically and physically bound to the substrate and (2) the high available “free water” where water can diffuse within the matrix to the microorganisms as well as organic substrates (Pommier and Chenu, 2007). When the total solids (TS) content increases, the quantity of free water decreases with a strong modification of the rheological behaviour of the substrate (García-Bernet et al., 2011). Thus, the transport of soluble compounds within the substrate can become a limiting factor (Bollon et al., 2013). Increasing the TS content usually leads to a reduction of hydrolysis and substrate degradation with a subsequent decrease in biogas production (Abbassi-Guendouz et al., 2012; García-Bernet et al., 2011; Motte et al., 2013). As an illustration, by increasing the TS content from 20 to 30%, Fernández et al. (2008) observed a decrease of organic waste degradation by 17%. At a TS higher than 30%, methane production was strongly inhibited by VFAs accumulation and decrease of pH (Abbassi-Guendouz et al., 2012).

Overall, the limitation of D-AD at high TS might be explained by the slow diffusion of dissolved molecules inside the organic matrix, generating a local accumulation of inhibitory products for methanogens at microbial scale (Martin, 2001; Staley et al., 2011). The inhibitory fermentative by-products could be not only Volatile Fatty Acids (VFAs) (Staley et al., 2011) but also dissolved

gases such as hydrogen (Abbassi-Guendouz et al., 2012). In a recent study, Abbassi-Guendouz et al. (2012) suggested that methanogenesis inhibition at solids content higher than 30% was likely caused by mass transfer limitation and more particularly by a local accumulation of fermentative gases, such as H<sub>2</sub> and CO<sub>2</sub>, leading to further VFAs accumulation.

The aim of this study was to investigate the inhibitory effect of dissolved gases on D-AD and especially H<sub>2</sub> and CO<sub>2</sub>, since they are both produced by acidogenesis. Batch experiments were designed and carried out to evaluate specifically the effect of H<sub>2</sub> and/or CO<sub>2</sub> partial pressure on methanogenesis in D-AD system.

## **MATERIAL AND METHODS**

Batch experiments were carried out in four replicates during 13 days, with wheat straw as substrate at 25% TS, pH fixed at 8, and substrate/inoculum ratio of 3 (Volatile Solids basis). After inoculation and moisture content adjustment, a thin layer (< 1 cm) of 1 mm meshed wheat straw was first placed at the bottom of the flasks to avoid the effect of gas diffusion within the matrix. Different initial partial pressure of gas (H<sub>2</sub>, CO<sub>2</sub> and H<sub>2</sub>/CO<sub>2</sub> mixture) were applied: from 0 to 1 530 mbars of H<sub>2</sub>, from 0 to 1 982 mbars of CO<sub>2</sub>, and H<sub>2</sub>/CO<sub>2</sub> mixture with H<sub>2</sub> and CO<sub>2</sub> ranging from 0 to 1 042 mbars and 0 to 433 mbars, respectively.

In a second part, another set of experiments was carried out by adding initially 1 000 mbars of H<sub>2</sub> and further addition of 400 mbars of CO<sub>2</sub> after different times of operation (11 and 18 days).

The medium was buffered and no pH variation was observed. The biogas was measured periodically throughout the experiments. The quantities of VFAs were measured at the start, before adding CO<sub>2</sub> and at the end of each experiment.


## **MAIN RESULTS**

### **Specific H<sub>2</sub> inhibition of methanogenesis and hydrolysis in D-AD**

From 0 to 600 mbars of initial H<sub>2</sub>, a slight increase of CH<sub>4</sub> production was observed from 25.2 ± 3.5 mL CH<sub>4</sub>/g<sub>VS</sub> in the controls to 39.6 ± 4.9 mL CH<sub>4</sub>/g<sub>VS</sub>. Such increase was due to the consumption of H<sub>2</sub> added initially. At higher initial p<sub>H<sub>2</sub></sub>, the cumulative production of CH<sub>4</sub> decreased from 39.6 ± 4.9 mL CH<sub>4</sub>/g<sub>VS</sub> for an initial p<sub>H<sub>2</sub></sub> of 600 mbars down to 8.45 ± 1.46 mL CH<sub>4</sub>/g<sub>VS</sub> for an initial p<sub>H<sub>2</sub></sub> of 1 555 mbars., with a transition phase between 650 and 800 mbars. Therefore, an initial p<sub>H<sub>2</sub></sub> higher than 800 mbars induced a strong and stable inhibition of methanogenesis at TS content of 25%.

No accumulation or significant change in VFAs was observed whatever the initial p<sub>H<sub>2</sub></sub>, and inhibition of the methanogenesis step was also not related to a pH decrease. In addition, the acetogenesis was not inhibited and no other metabolic pathways were found for a high p<sub>H<sub>2</sub></sub> like for example a production of butyrate or caproate (Arslan et al., 2012). Therefore, VFA accumulation or changes in pattern cannot explain the observed methanogenesis inhibition, as previously shown in wet AD (Ahring and Westermann, 1988; Amani et al., 2010; Siegert and Banks, 2005).

When considering the total amount of metabolic products (VFAs, H<sub>2</sub>), the total substrate degradation rate (Figure 1) decreased according to the initial p<sub>H<sub>2</sub></sub>. Since no sugars were accumulated in the bulk phase, these results suggest a strong inhibition of hydrolysis at initial partial pressure of H<sub>2</sub> higher than 800 mbars. It is well known that hydrolysis is the limiting step in D-AD (Pavlostathis and Giraldo-Gomez, 1991) and the composition of the substrate (morphological and chemical) can have a strong effect on this step, especially the quantity of lignin for lignocellulosic wastes (Monlau et al., 2012). The present study clearly shows for the first time a strong inhibition of hydrolysis by H<sub>2</sub> partial pressure in D-AD.



**Figure 1:** Substrate degradation ( g COD at the end/g TS –g COD at the beginning/g TS) according to the initial partial pressure of H<sub>2</sub> in mbars at 25% of TS, 35°C and pH 8. TS: Total Solids; COD: Chemical oxygen demand.

Interestingly, at low initial  $p_{H_2}$ , H<sub>2</sub> was totally converted to acetate and methane. At an initial  $p_{H_2}$  higher than 750 mbars, H<sub>2</sub> was not fully consumed and H<sub>2</sub> remained in headspace at the end of the experiments. From 750 to 1 000 mbars, a part of initial H<sub>2</sub> was used to produce acetate by homoacetogenesis (Montero et al., 2009) but H<sub>2</sub> was not totally consumed since some H<sub>2</sub> remained at the end of the experiment. CO<sub>2</sub> accumulation was also monitored and at a  $p_{H_2}$  ranging from 0 to 750 mbars, CO<sub>2</sub> was produced due to the microbial activity and a decrease of the released CO<sub>2</sub> was observed from  $38.98 \pm 10.45$  to  $15.33 \pm 2.3$  ml CO<sub>2</sub> /g<sub>VS</sub> in the controls and at 750 mbars of  $p_{H_2}$  respectively. Over 750 mbars, no CO<sub>2</sub> was accumulated in the headspace. Since the increase of the remaining amount of H<sub>2</sub> in headspace was correlated to a decrease of remaining CO<sub>2</sub>, it was concluded that CO<sub>2</sub> production was the limiting factor for further H<sub>2</sub> consumption (Florin and Harris, 2008). This hypothesis was tested by adding both CO<sub>2</sub> (400 mbars) and H<sub>2</sub> (1 000 mbars) at the start of the experiment. After correcting methane production by the methane produced using the added gases, methane production was similar in the mixture of H<sub>2</sub>/CO<sub>2</sub> and in the control. Therefore, in that case, no methanogenesis inhibition occurred. These results confirm that CO<sub>2</sub> production can be considered as the limiting factor when  $p_{H_2}$  is high.

### Reversibility of the H<sub>2</sub> inhibition in D-AD

Since adding CO<sub>2</sub> can prevent the H<sub>2</sub> inhibition of hydrolysis and methanogenesis, we investigated the reversibility of the H<sub>2</sub> inhibition by introducing CO<sub>2</sub>. An inhibitory  $p_{H_2}$  of 1 000 mbars was added at the start of the experiment and CO<sub>2</sub> (400 mbars) was further added at different times (11 and 18 days) to test the reversibility of the inhibition.

For both experiments, H<sub>2</sub> inhibition stopped after CO<sub>2</sub> addition. The profiles of cumulated production of CH<sub>4</sub> after adding the CO<sub>2</sub> shows: (1) a rapid increase of CH<sub>4</sub> production mainly due to H<sub>2</sub> and CO<sub>2</sub> consumption, (2) a short lag phase with a slower production of CH<sub>4</sub> and (3) a constant and stable CH<sub>4</sub> production and substrate degradation as observed in the control. These last

experiments clearly showed the reversibility of hydrolysis inhibition by H<sub>2</sub>.

## CONCLUSION

This study provides new insight about the effect of fermentative gases and more particularly H<sub>2</sub>, on D-AD when using wheat straw as substrate. By adding CO<sub>2</sub>, hydrolysis inhibition by H<sub>2</sub> was not only suppressed but was also fully reversible. This indicates that CO<sub>2</sub> could be a limiting factor in D-AD when inhibition by H<sub>2</sub> occurs. Our results were however obtained with a thin layer substrate to avoid the effect of the gas diffusion. In real D-AD digesters, combined effects of inhibition as well as gas diffusion can occur. Nevertheless, hydrolysis inhibitory effect of H<sub>2</sub> can take place at a local scale, and may be one of the main reasons of the decrease in overall digester performances in D-AD.

## ACKNOWLEDGEMENTS

This work was supported by the French Environment and Energy Management Agency (ADEME) and the French Institute for Agricultural and Food Research (INRA).

## REFERENCES

- Abbassi-Guendouz, A., Brockmann, D., Trably, E., Dumas, C., Delgenès, J.-P., Steyer, J.-P., Escudié, R., 2012. Total solids content drives high solid anaerobic digestion via mass transfer limitation. *Bioresour. Technol.* 111, 55–61.
- Ahring, B.K., Westermann, P., 1988. Product inhibition of butyrate metabolism by acetate and hydrogen in a thermophilic coculture. *Appl. Environ. Microbiol.* 54, 2393–7.
- Arslan, D., Steinbusch, K.J.J., Diels, L., De Wever, H., Buisman, C.J.N., Hamelers, H.V.M., 2012. Effect of hydrogen and carbon dioxide on carboxylic acids patterns in mixed culture fermentation. *Bioresour. Technol.* 118, 227–34.
- Bollon, J., Benbelkacem, H., Gourdon, R., Buffière, P., 2013. Measurement of diffusion coefficients in dry anaerobic digestion media. *Chem. Eng. Sci.* 89, 115–119.
- Florin, N.H., Harris, A.T., 2008. Enhanced hydrogen production from biomass with in situ carbon dioxide capture using calcium oxide sorbents. *Chem. Eng. Sci.* 63, 287–316.
- García-Bernet, D., Buffière, P., Latrille, E., Steyer, J.-P., Escudié, R., 2011. Water distribution in biowastes and digestates of dry anaerobic digestion technology. *Chem. Eng. J.* 172, 924–928.
- Martin, D.J., 2001. The site of reaction in solid-state digestion: a new hypothesis. *Trans IChemE* 79, part B.
- Monlau, F., Sambusiti, C., Barakat, A., Guo, X.M., Latrille, E., Trably, E., Steyer, J.P., Carrere, H., 2012. Predictive models of biohydrogen and biomethane production based on the compositional and structural features of lignocellulosic materials. *Environ. Sci. Technol.* 46, 12217–12225.
- Montero, B., Garcia-Morales, J.L., Sales, D., Solera, R., 2009. Analysis of methanogenic activity in a thermophilic-dry anaerobic reactor: use of fluorescent in situ hybridization. *Waste Manag.* 29, 1144–51.
- Motte, J.-C., Escudié, R., Bernet, N., Delgenes, J.-P., Steyer, J.-P., Dumas, C., 2013. Dynamic effect of total solid content, low substrate/inoculum ratio and particle size on solid-state anaerobic digestion. *Bioresour. Technol.* 144, 141–148.
- Pavlostathis, S.G., Giraldo-Gomez, G., 1991. Kinetics of anaerobic treatment. *Water Sci. Technol.* 24, 35–59.
- Pommier, S., Chenu, D., 2007. A logistic model for the prediction of the influence of water on the solid waste methanization in landfills. *Biotechnol. Bioeng.* 97, 473–482.
- Staley, B.F., de Los Reyes, F.L., Barlaz, M. a, 2011. Effect of spatial differences in microbial activity, pH, and substrate levels on methanogenesis initiation in refuse. *Appl. Environ. Microbiol.* 77, 2381–2391.