

The Lime Leaf Miner Phyllonorycter issikii (Kumata, 1963), a Highly Invasive Pest in Europe: Genetics of Invasion and Systematic

Natalia Kirichenko, Paolo Triberti, Sylvie Augustin, Alain Roques, Carlos Lopez-Vaamonde

▶ To cite this version:

Natalia Kirichenko, Paolo Triberti, Sylvie Augustin, Alain Roques, Carlos Lopez-Vaamonde. The Lime Leaf Miner Phyllonorycter issikii (Kumata, 1963), a Highly Invasive Pest in Europe: Genetics of Invasion and Systematic. 8. International Conference on Biological Invasions from understanding to action "NEOBIOTA 2014", Nov 2014, Antalya, Turkey. 279 p. hal-02740096

HAL Id: hal-02740096 https://hal.inrae.fr/hal-02740096

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The Lime Leaf Miner *Phyllonorycter issikii* (Kumata, 1963), a Highly Invasive Pest in Europe: Genetics of Invasion and Systematic

Natalia KIRICHENKO^{1,2}, Paolo TRIBERTI³, Sylvie AUGUSTIN² Alain ROQUES², Carlos Lopez-VAAMONDE²

¹V.N. Sukachev Institute of Forest SB RAS, Krasnoyarsk, 660036 Russia Email: nkirichenko@yahoo.com

Herbivore insects are one the most numerous invaders at many locations around the world, representing an economic threat, causing serious impacts on communities of native species, and disturbing natural ecosystem processes. Among those invasive plant-feeding pests, leaf mining insects (larvae developed within leaves) represent an important group of herbivores that threaten crops, parks and gardens. Some leaf mining micromoths provide extraordinary examples of rapid expansion, particularly the lime leaf miner *Phyllonorycter issikii*, a *Tilia*-feeder. A native of Eastern Asia, in the last few decades, this tiny moth has spread westwards over the whole of Russia and has invaded several European countries, becoming a serious ornamental pest of lime trees.

Within a project supported by LE STUDIUM® Loire Valley Institute for Advanced Studies (France), we are analyzing the genetics of the invasion of *P. issikii* across its distribution range using mitochondrial (COI) and nuclear (28S and Histone3) gene fragments. In addition, the morphology of barcoded adult moths is being compared between native and invaded areas. Preliminary results indicate a loss of genetic diversity in *P. issikii* populations, following the range spread from East to West. Both genetic and morphometric data support the existence of a new undescribed non-invasive cryptic species that occurs sympatrically with *P. issikii* in the Russian Far East.

Keywords: Phyllonorycter issikii, invasion, Europe, genetics, systematics

²INRA Institut National de la Recherche Agronomique, URZF, Orléans cedex 2, F-45075, France

³Museo Civico di Storia Naturale, Verona, I37129, Italy