



**HAL**  
open science

## Resilience and resistance of microbial communities against drought events, in a mature agroforestry system in Mediterranean area

Esther Guillot, Philippe Hinsinger, Lydie Dufour, Isabelle Bertrand

### ► To cite this version:

Esther Guillot, Philippe Hinsinger, Lydie Dufour, Isabelle Bertrand. Resilience and resistance of microbial communities against drought events, in a mature agroforestry system in Mediterranean area. 5. International EcoSummit: Ecological Sustainability: Engineering Change, Aug 2016, Montpellier, France. 2016. hal-02740136

**HAL Id: hal-02740136**

**<https://hal.inrae.fr/hal-02740136>**

Submitted on 2 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License



# Resilience and resistance of microbial communities against drought events, in a mature agroforestry system in Mediterranean area



Guillot Esther<sup>1</sup>, Dufour Lydie<sup>2</sup>, Hinsinger Philippe<sup>1</sup>, Bertrand Isabelle<sup>1</sup>

1. UMR Eco&Sols, INRA, Montpellier, France




2. UMR System, INRA, Montpellier, France

esther.guillot@gmail.com



- Agroforestry systems are known to limit soil degradation, deeply store carbon and have positive impact on different indicators of soil quality.
- What is the impact of tree-crop associations on resistance and resilience of topsoil microbial communities?
- Our aim is to evaluate the resistance and the resilience of the microbial communities of topsoil along a spatial gradient from the tree line to the center of the intercrop, and in the monocrop treatment.

## Agroforestry system



	Day	Distance				
		0-1m	1-2m	2-4m	4-6.5m	Monocrop treatment
Rate day <sup>-1</sup> µg C-CO <sub>2</sub> g <sup>-1</sup> dry soil	Day 21 : Resistance	1.315 <sup>a</sup>	1.579 <sup>a</sup>	1.823 <sup>a</sup>	0.780 <sup>a</sup>	1.543 <sup>a</sup>
	Day 29 : Resilience	18.556 <sup>a</sup>	14.670 <sup>ab</sup>	9.125 <sup>b</sup>	12.579 <sup>ab</sup>	7.503 <sup>ab</sup>
Index	Day 21 : Resistance	0.0758 <sup>a</sup>	0.147 <sup>a</sup>	0.373 <sup>a</sup>	0.122 <sup>a</sup>	0.357 <sup>a</sup>
	Day 29 : Resilience	0.216 <sup>ab</sup>	0.607 <sup>a</sup>	0.219 <sup>ab</sup>	0.477 <sup>ab</sup>	-0.064 <sup>b</sup>

Table 1 : Daily rate of C mineralisation at the end of the stress (Resistance) and 7 days after rewetting (Resilience) and indices of resistance and resilience based on cumulated C mineralisation. Data are mean, n=5, letters indicate significant differences (P<0.05)

- Stronger cumulated C mineralisation occurring close to the tree, decreasing with increasing distance from tree.
- Our data show difference in microbial resilience along the spatial gradient, but not difference in resistance.
- Microbial communities seem to be more sensitive to water stress and rewetting close to the tree than in the monocrop treatment.