



# Grassland Fabaceae grown under contrasted phosphorus supply induced changes in rhizospheric soil phosphatase activity

Florian Fort, Pablo Cruz, Stroia Ciprian, Claire Jouany

## ► To cite this version:

Florian Fort, Pablo Cruz, Stroia Ciprian, Claire Jouany. Grassland Fabaceae grown under contrasted phosphorus supply induced changes in rhizospheric soil phosphatase activity. Phosphorus in Soils and Plants: 5. International Symposium, Aug 2014, Montpellier, France. 148 p., 2014, PSP5 2014. Facing Phosphorus Scarcity. hal-02740155

**HAL Id: hal-02740155**

**<https://hal.inrae.fr/hal-02740155>**

Submitted on 2 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

> Le Corum, Montpellier, France

> **26-29 August 2014**

**PSP5 2014**

*Facing Phosphorus Scarcity*

# Phosphorus in Soils and Plants





**Theme 1 – Phosphorus forms, availability and cycling in soils**



## **Theme 1 – Phosphorus forms, availability and cycling in soils**

### **Keynote presentation**

## P207

### Grassland Fabaceae grown under contrasted phosphorus supply induced changes in rhizospheric soil phosphatase activity

Florian Fort <sup>1</sup>, Pablo Cruz <sup>1</sup>, Ciprian Stroia <sup>2</sup>, Claire Jouany <sup>1</sup>

1. INRA, AGIR UMR 1248, 24 Chemin de Borde Rouge – Auzeville CS 52627, 31326 Castanet Tolosan cedex, France

2. Banat University of Agricultural Sciences and Veterinary Medicine, Department of Biology and Plant Protection, Calea Aradului 119, 300645 Timisoara, Romania

Fabaceae performances in low fertility agro systems rely on their capacity to acquire phosphorus (P) since their growth is highly limited by P shortage. Recent work conducted on grassland's legumes demonstrated, that different roots strategies exist within fabaceae family resulting in a large range of responses to P stress. Our objective is to evaluate to what extent rhizospheric soil phosphatase activity is related to Fabaceae response to P limitation. In that purpose, we grew in a green house, 13 grassland Fabaceae species under two levels of P availability for more than 100 days. At harvest, rhizospheric soil were sampled then analysed for alkaline phospho-monoesterase activity together with bulk soil (control). Results show a significant effect of plant on species identity on rhizospheric phosphatase activity, *Anthyllis vulneraria* and *Vicia cracca* induce high phosphatase activity in their rhizospheric soil while *Securigera varia* and *Trifolium pratense* induce low phosphatase activity. There is also a significant and negative effect of P supply on phosphatase activity of rhizospheric soil. The activity measured for the bulk soil is always significantly lower than the one measured for the rhizospheric soil. It was interesting to notice that phosphatase activities measured for Fabaceae are not different from those obtained for Poaceae grown under similar conditions. We showed that Fabaceae species induce a strong increase of the phosphatase activity in their rhizospheric soil in comparison to bulk soil. However, further work is needed to understand the link between rhizospheric soil activity and P stress tolerance.

## P208

### A simple technique for studying root hair development under varying levels of buffered P supply

Elke Vandamme <sup>1</sup>, Marian Renkens <sup>2</sup>, Erik Smolders <sup>2</sup>, Roel Merckx <sup>2</sup>

1. Africa Rice Center (AfricaRice), Mikocheni B/Kawe, Avocado Street, P.O. Box 33581, Dar es Salaam, Tanzania

2. Division Soil and Water Management, Department of Earth and Environmental Sciences, KU Leuven, Kasteelpark Arenberg 20, 3001 Leuven, Belgium

Root hairs contribute significantly to phosphorus (P) uptake and are particularly important when P supply is low and strongly diffusion-limited, which is the case for many depleted and P-fixing soils in the tropics. In view of the need to select and develop P-efficient crops, it may be of interest to explore genotypic variation in root hair development. However, the measurement of root hair development of soil-grown plants is difficult as root hairs are easily damaged while roots are washed out from soil. Growth pouches or hydroponics have been used to study root hair development but in these systems it is not possible to spatially buffer P supply. A method was developed to evaluate root hair growth under artificial conditions that closely mimic diffusion-limited P supply in soils. The method involves seedlings grown on an agar medium that contains P adsorbed on Al<sub>2</sub>O<sub>3</sub> nanoparticles at different levels of P intensity. The use of Al<sub>2</sub>O<sub>3</sub> nanoparticles as P buffer ensures clarity of the agar, allowing for in-situ measurements of root hair length and density. Results of a study on 8 soybean genotypes screened in both the agar medium and a pot experiment using a P-deficient soil at low and high P supply are presented. P intensity in the agar was buffered at 6 and 80 µM, while total P concentrations in the medium were 120 and 250 µM respectively. The P intensities corresponded to the P concentrations measured in the pore water of the amended soil samples used for the plant growth experiment. We conclude that the technique provides a simple way to study the plastic response of root hairs to P deficiency governed by localized P-sensing mechanisms.