



Regional control over B concentration and $\delta^{11}\text{B}$ atmospheric input

Philippe Roux, Marie-Pierre M.-P. Turpault, Paul Olivier Redon, Damien
Lemarchand

► To cite this version:

Philippe Roux, Marie-Pierre M.-P. Turpault, Paul Olivier Redon, Damien Lemarchand. Regional control over B concentration and $\delta^{11}\text{B}$ atmospheric input. 25. Goldschmidt Conference, Aug 2015, Prague, Czech Republic. hal-02740305

HAL Id: hal-02740305

<https://hal.inrae.fr/hal-02740305>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Regional control over B concentration and $\delta^{11}\text{B}$ atmospheric input

P. ROUX^{*12}, M-P. TURPAULT¹, P-O. REDON³ AND D. LEMARCHAND²

¹BEF-INRA, Centre de Nancy, 54280 Champenoux, France
(*correspondence : philippe.roux@nancy.inra.fr)

²LHyGeS/CNRS, Université de Strasbourg, France
(lemarcha@unistra.fr)

³Andra, Centre de Meuse/Haute-Marne, 55290 Bure, France

Boron concentration and isotopic composition ($\delta^{11}\text{B}$) of atmospheric inputs (openfield dust and total dissolved depositions) were studied for the first time over a two-year period (2012-2013) to determine the budgets, sources and processes controlling the atmospheric contribution of boron in a forest ecosystem. Boron concentrations in dust depositions (Du) show very high B concentrations ranging from $191 \mu\text{g.g}^{-1}$ to $501 \mu\text{g.g}^{-1}$ while most of total dissolved depositions (TDi) ranges from 1.1 ng.mL^{-1} to 3.6 ng.mL^{-1} with an extreme event at 9.1 ppb during the summer of 2013. $\delta^{11}\text{B}$ show a large range of variations, from -4.7 ‰ to +11.4 ‰ for Du and from 8.3 ‰ to 36.6 ‰ for TDi. The measured concentrations yield an annual boron flux of about $10 \text{ mg.ha}^{-1}.\text{yr}^{-1}$ for Du and about $16 \text{ mg.ha}^{-1}.\text{yr}^{-1}$ for TDi.

Du data show no clear relationship between B and major or trace elements but the unexpected and rather constant high B concentrations observed over the sampling period indicate a control by one or more B-rich sources, yet to be determined. The $\delta^{11}\text{B}$ vs. $1/\text{B}$ mixing diagram evidences the contribution of at least three components. B analyses of some of the fertilizers spread in the nearby agricultural soils and samples of top soil fine particules will allow testing the hypothesis of a regional origin of Du in relation with agricultural activities.

$\delta^{11}\text{B}$ data in TDi show a clear seasonal cyclicity throughout the sampling period and are characterized by a high isotopic signature during the summer and a low isotopic signature during the winter. However, the B concentrations remain roughly stable without clear seasonal cyclicity. The observed high isotopic ratios are not associated with a high seawater contribution (according to Na) as classically proposed. This indicates that marine-derived particles unlikely control dissolved boron in our ecosystem. Furthermore, no clear relationship is observed between B, $\delta^{11}\text{B}$ and the dissolved major elements, evidencing no major source contribution. We hypothesize that dissolved boron mostly originates from condensated gaseous boron. Evaporation/condensation processes appear therefore to be controlled by local physical atmospheric parameters.