The Changing Face of Septic Arthritis Complicating Rheumatoid Arthritis in the Era of Biotherapies. Retrospective Single-Center Study over 35 Years

Jean Jacques Dubost, Bruno Pereira, Anne Tournadre, Zuzana Tatar, Marion Couderc, Martin Soubrier

► To cite this version:

Jean Jacques Dubost, Bruno Pereira, Anne Tournadre, Zuzana Tatar, Marion Couderc, et al.. The Changing Face of Septic Arthritis Complicating Rheumatoid Arthritis in the Era of Biotherapies. Retrospective Single-Center Study over 35 Years. 2015 ACR/ARHP Annual Meeting, Nov 2015, San Francisco, United States. 4053 p., 10.1002/art.39448 . hal-02740404

HAL Id: hal-02740404

<https://hal.inrae.fr/hal-02740404>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution| 4.0 International License

The Changing Face of Septic Arthritis Complicating Rheumatoid Arthritis in the Era of Biotherapies. Retrospective Single-Center Study over 35 Years

Jean-Jacques Dubost¹, Bruno Pereira², Anne Tournadre³, Zuzana Tatar¹, Marion Couderc¹ and Martin Soubrier¹, ¹Rheumatology department CHU Clermont-Ferrand, Clermont-Ferrand, France, ²Biostatistics unit (DRCI), CHU Clermont-Ferrand, Clermont-Ferrand, France, ³Rheumatology, UNH-UMR 1019 INRA University of Auvergne and Rheumatology department CHU Clermont-Ferrand, Clermont-Ferrand, France

Meeting: **2015 ACR/ARHP Annual Meeting**

Date of first publication: September 29, 2015

Keywords: **Arthritis, Biologic drugs, infection and rheumatoid arthritis (RA)**

SESSION INFORMATION

Date: **Monday, November 9, 2015**

Session Type: ACR Poster Session B

Session Title: **Rheumatoid Arthritis - Clinical Aspects Poster II**

Session Time: 9:00AM-11:00AM

Background/Purpose: Rheumatoid arthritis (RA) is a risk factor for septic arthritis (SA), and anti-TNF therapy doubles the risk of SA. The purpose of this study was to see whether the frequency and features of SA complicating RA have changed over the last 35 years.

Methods: This retrospective single-center study included the full register of all patients hospitalized at the rheumatology department of Clermont-Ferrand—CHU between 1979 and 2013 for septic arthritis bacteriologically documented by synovial fluid and/or blood culture samples. The periods 1979–2002 (period before biotherapies) and 2003–2013 (last decade in the era of biotherapies) were compared.

Results: Between 1979 and 2013, 64/514 (12.5%) SA presented with a RA—21/153 (13.7%) in the 2003–2013 period and 43/361 (11.9%) in the 1979–2002 period. Over the past decade, median age of SA-RA patients increased (61 vs 68 years; $p<0.02$) and predominant gender became males (52% vs 40%). The features of the RA remained unchanged: history (18 years (8–29) vs 16 (8–25)), rheumatoid factor (95% vs 87%), and corticosteroids (91% vs 81%) at the same mean dose (10 mg/d). 71% in the period before biotherapies and 63% in the last decade received a DMARD. Over the last decade 24% (vs 0; $p<0.003$) of patients received a biologic DMARD: etanercept ($n=2$), adalimumab ($n=1$), rituximab ($n=1$), tocilizumab ($n=1$). Proportion of polyarticular infection had decreased (9.5% vs 37%; $p<0.02$), down to the same level as SA-non-RA cases (8%). Proportion of *S. aureus* infections had stabilized (62% vs 74%) in SA-RA patients but was higher than SA-non-RA (47% and 53%). MRSA infections became more frequent in SA-RA (31% vs 6%; $p<0.05$) in contrast to SA-non-RA cases (8% vs 16%; ns). Gram-negative bacilli infections have tended to become more frequent (19% vs 5%; $p=0.08$). Blood cultures less often tested positive (29% vs 47%; ns). Mortality rates has fallen slightly (5% vs 9%; ns), in contrast to SA-non-RA cases (7% vs 6%; ns).

Conclusion: This study brings reassuring findings—in the era of biotherapies, the frequency of septic arthritis complicating rheumatoid arthritis has stabilized, and the most severe septic polyarticular forms are on the decline.

Disclosure: J. J. Dubost, None; B. Pereira, None; A. Tournadre, None; Z. Tatar, None; M. Couderc, None; M. Soubrier, None.

To cite this abstract in AMA style:

Dubost JJ, Pereira B, Tournadre A, Tatar Z, Couderc M, Soubrier M. The Changing Face of Septic Arthritis Complicating Rheumatoid Arthritis in the Era of Biotherapies. Retrospective Single-Center Study over 35 Years [abstract]. *Arthritis Rheumatol*. 2015; 67 (suppl 10). <http://acrabstracts.org/abstract/the-changing-face-of-septic-arthritis-complicating-rheumatoid-arthritis-in-the-era-of-biotherapies-retrospective-single-center-study-over-35-years/>. Accessed April 5, 2016.

ACR Meeting Abstracts - <http://acrabstracts.org/abstract/the-changing-face-of-septic-arthritis-complicating-rheumatoid-arthritis-in-the-era-of-biotherapies-retrospective-single-center-study-over-35-years/>