



Quantitative relationships between the tissue composition of bovine carcass and easily obtainable indicators

Marwa Al-Jammas, Jacques Agabriel, Jean Vernet, Isabelle Ortigues Marty

► To cite this version:

Marwa Al-Jammas, Jacques Agabriel, Jean Vernet, Isabelle Ortigues Marty. Quantitative relationships between the tissue composition of bovine carcass and easily obtainable indicators. 61. International Congress of Meat Science and Technology (ICoMST), Aug 2015, clermont-Ferrand, France. hal-02740472

HAL Id: hal-02740472

<https://hal.inrae.fr/hal-02740472>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

QUANTITATIVE RELATIONSHIPS BETWEEN THE TISSUE COMPOSITION OF BOVINE CARCASS AND EASILY OBTAINABLE INDICATORS

M. AL-JAMMAS^{1,2}, J. AGABRIEL^{1,2}, J. VERNET^{1,2} and I. ORTIGUES-MARTY^{1,2}

¹ INRA, UMR1213 Herbivores, F-63122 Saint-Genès-Champanelle, France

² Clermont Université, VetAgro Sup, UMR1213 Herbivores, BP 10448, F-63000, Clermont-Ferrand, France

Abstract – The most accurate determination of beef carcass quality involves dissection of cut or entire carcass. This, however, is very costly and cumbersome. An alternative is to determine easily obtainable indicators. The objective of this study was to derive quantitative relationships between indirect carcass indicators and measured carcass tissue composition. A meta-analysis was applied on 25 published trials with cattle. The selected indicators were USDA yield grade, fat thickness, marbling, ribeye area and carcass conformation and fatness scores. The USDA yield grade was the most highly related to changes in carcass adipose tissue and muscle weights. Fat thickness was also related to changes in both adipose tissue and muscle weights. Other indicators were less correlated with changes in carcass tissue composition. The relationships obtained in this study depend on measurement accuracy and different deposition kinetics for adipose tissue.

Key Words – adipose tissue, carcass composition, cattle.

I. INTRODUCTION

Carcass quality depends on the tissue composition and the muscularity of slaughtered animals. Information about carcass composition helps to determine nutritional requirement, animal performance and the cost of production of beef cattle. Carcass quality in cattle is evaluated by a variety of methods, each of which may involve different techniques. Direct methods determine the composition (tissue or chemical) of the carcass from the complete dissection of the half or the entire carcass. The most accurate method is the chemical analysis of the carcass [1]. However, the physical separation of the carcass into muscle, adipose tissue and bone is a useful indicator of retail composition [2] or tissue development [3]. These measures are difficult and expensive to implement in practice particularly with cattle.

Indirect methods are commonly used. In Europe the conformation (ConS) and fatness (FS) scores are used. In other countries as USA, the yield grade (YG), subcutaneous fat thickness (SFT), marbling (Mar) and ribeye area (longissimus dorsi area, LMA) are used. Therefore, the comparison of published quantitative results on carcass quality is difficult because of this diversity of approaches. We recently showed [4] that variations in yield grade and fat thickness may properly explain the variations of chemical composition of carcass. The objective of this study was to determine, from published results, the quantitative relationships between variations of tissue composition of carcass and the variation of easily obtainable indicators.

II. MATERIALS AND METHODS

Publications describing cattle carcass and published in international scientific journals were searched using the web of science without restriction on the years. Selection criteria were results of carcass tissue composition with at least one indicator (YG, SFT, LMA, Mar, Cons and FS). Data extracted from the publications included detailed descriptions of animals (breed, gender...etc) and carcass characteristics. They were entered into a database.

Several additional calculations were performed to complete the database in case of missing data. They were based on data available in the publication.

Cold carcass weight, kg = $0.98 \times$ hot carcass weight (HCW, kg).

Dressing percentage, % = $(\text{hot carcass weight} \div \text{slaughter weight}) \times 100$

The marbling score was harmonized on a scale of 10 (2: traces, 3: slight, 4: small) and the conformation and fatness scores were harmonized on scales of 5 [5].

Publications presenting more than 1 study were separated into experiments that were individually encoded as such. All data were examined graphically in several steps during the analysis. Descriptive statistics (mean, standard deviation, minimum, maximum) and the normality test for each variable were generated. To take in account the other factors of variation (gender, breeds...) methods of meta-analysis were applied in intra-study according to *Sauvant et al.* [6]. Variance-covariance models were developed using Minitab software (Minitab® 16.2.4. 2013), as follows:

$$Y_{ij} = \mu + \mu_i + \beta X_{ij} + \beta_i X_{ij} + e_{ij}$$

Where, i : the index of the study factor, j the index of the treatment number, Y_{ij} the dependent variable, X_{ij} : the quantitative explanatory variable, μ the intercept, μ_i the fixed effect of the study on the intercept, β : the general fixed regression coefficient, β_i the fixed effect of the study on slope, and e_{ij} the unexplained residual error.

The response variable was the tissue composition of carcass (muscle, adipose tissues in kg). The independent variables for each relationship were the hot carcass weight in kg, associated with one carcass indicator (yield grade, fat thickness in mm, marbling score /10, ribeye area in cm², conformation score /5 and fatness score/5). For all models, the normality of residuals was verified, and influent data were identified from analysis of residuals, high leverage, Cook's distance and DEFIT [6]. The contribution of each explanatory variable to the total variance of the model was calculated. The models were compared in terms of adjusted R², residual means square errors of the model (RMSE), and the mean square error of prediction (MSPE) [7]:

$MSPE = 1/n \sum_{i=1}^n (X_i - y_i)^2$, where x are predicted values; y are observed values.

III. RESULTS AND DISCUSSION

A total of 25 scientific papers were identified, they included 28 experiments for a total of 117 treatments. The animals differed in gender (4% female, 43% male, 49% castrated males and 4% implanted castrated males), and/or breeds (45% beef, 29% dual purpose and 26% dairy). The descriptive statistics for variables retrieved from

the publications showed wide ranges in all live animal and carcass characteristics (Table 1). The majority of data (88%) were associated with the continental breeds or their cross breeds with the lowest adipose tissue weights and the highest HCW and muscle weights. The remaining 12 % of data came from British breeds. Overall HCW varied between 138 - 487 kg, with 67% of data between 300-380 kg. Adipose tissues varied between 11-137 kg, with 81% of data between 30-80 kg, representing 16-24% of HCW.

Table 1: Mean values, standard deviations (SD), minimum (Min) and maximum (Max) of carcass weight, tissue composition and indicators.

Variable	Number ¹		Mean	SD	Min	Max
	n _e	n _m				
Hot carcass weight, kg	116	1	313.8	63.6	138.3	487.5
Cold carcass weight, kg	116	1	307.5	62.4	135.5	477.8
Adipose tissue, kg	116	1	51.6	27.2	11.1	137.2
Muscle, kg	116	1	205.3	54.9	93.1	364.1
Bone, kg	116	1	51.7	12.8	24.6	76.5
Fat thickness, mm	60	57	8.8	3.9	3.2	20.2
Marbling /10	43	74	3.8	0.9	2.3	6.3
Yield grade	22	95	2.7	0.9	1.1	4.7
Ribeye area, cm ²	79	38	68.3	13.6	38.9	106.0
Conformation score /5	67	50	2.9	0.8	0.7	4.8
Fatness score /5	72	45	2.9	0.8	1.4	4.8

1: number of data (treatment): n_e: number of existing data, n_m: number of missing data.

Relationships between carcass tissue composition and indicators are presented in Table 2. Hot carcass weight was more highly correlated with the muscle weight than with adipose tissue weight (Model 7 vs. 1) explaining a greater proportion of the variance (74 vs. 3%). Relationships were generally improved when indicators were included as additional covariate. Most indicators (YG, SFT, Mar, FS,) were positively correlated with adipose tissue weight (model 2, 3, 4, 6) and negatively correlated with muscle weight (Models 8, 9, 10, 13), contrary to LMA and ConS (Model 5, 11). These results can be explained by the negative correlations between adipose tissue and muscle weights at similar carcass weight.

Table 2. Intra-study relationships between carcass tissue weights (adipose tissue and Muscle,kg) and easily obtainable indicators (yield grade, subcutaneous fat thickness, marbling score, longissimus dorsi area, and score of conformation and fatness of carcass) in finishing cattle.

N°	Y	Number ¹		Equation ²	RMSE	Adj-R ²	MSEP	% variance explained ³		
		N _{sy}	N _{tr}					In	HCW	Sy
1	Adipose tissue, kg	28	117	24.41±9.26 ^{***} + 0.10±0.03 ^{***} HCW	9.05	0.89	61.5	-	3	88
2		7	22	-10.71±11.8 ^{NS} + 0.12±0.04 ^{***} HCW+20.00±2.99 ^{***} YG	5.74	0.98	19.5	62	26	10
3		14	57	2.08±8.98 ^{NS} + 0.12±0.03 ^{***} HCW +3.51±0.35 ^{***} SFT	4.65	0.98	15.6	74	2	22
4		11	43	3.52± 10.03 ^{NS} +0.09±0.04 [*] HCW +12.28±1.37 ^{***} Mar	4.91	0.97	16.9	24	35	38
5		16	76	58.44±11.05 ^{***} +0.22±0.04 ^{***} HCW -0.81±0.20 ^{***} LMA	8.99	0.91	61.8	5	15	73
6		15	72	-1.77±8.28 ^{NS} +0.04±0.02 [*] HCW +12.41±1.68 ^{***} FS/5	5.55	0.90	23.5	17	0.1	75
7	Muscle, kg	28	117	-38.54±9.27 ^{***} + 0.76±0.02 ^{***} HCW	9.06	0.97	61.5	-	74	24
8		7	22	1.04±7.07 ^{NS} + 0.71±0.03 ^{***} HCW -18.04±1.79 ^{***} YG	3.44	0.99	7.0	0	89	10
9		14	57	-11.61±6.20 ^{NS} +0.70±0.02 ^{***} HCW -2.92±0.24 ^{***} SFT	3.21	0.99	7.4	5	82	13
10		11	43	-9.62±9.43 ^{NS} +0.70±0.04 ^{***} HCW -9.17±1.29 ^{***} Mar	4.62	0.99	14.9	0	70	1
11		16	76	-73.69±10.83 ^{***} +0.62±0.04 ^{***} HCW +0.94±0.20 ^{***} LMA	8.82	0.97	59.3	1	69	27
12		14	67	-54.88±10.87 ^{***} +0.79±0.03 ^{***} HCW +5.55±2.87 [*] ConS	8.18	0.97	50.9	21	64	13
13		15	72	-15.11±8.62 [*] +0.83±0.02 ^{***} HCW -12.93±1.74 ^{***} FS	5.78	0.98	25.5	13	72	13

1: number of data: Nsy: number of studies, Ntr: number of treatments.

2: YG: Yield grade, HCW: Hot carcass weight, SFT: subcutaneous fat thickness, Mar: Marbling, LMA: Longissimus area, ConS: conformation score, FS: fatness score..

3: In: indicator (YG, SFT, Mar, LMA, ConS and FcS), Sy: Study.

NS: not significant $P>0.10$, *: $P<0.05$, **: $P<0.01$, ***: $P<0.001$.

The SFT and YG were the most highly related to changes of carcass adipose tissue weight (Model 1, 2) with the highest adjusted R^2 and among the lowest RMSE and MSEP. The proportions of variance explained by the study effect were the lowest, indicating that there were no major additional variation factors besides the 2 covariates. These two indicators were already shown to be well related to the proportion of lipids in the carcass [4]. Besides, SFT is considered in the calculation of yield grade; it expresses the quantity of subcutaneous adipose tissue which is related positively to total adipose tissue between birth and adult age [8]. Marbling, despite its relative subjectivity [9] was also related to carcass adipose tissue, even though the

quality of the relationship was lower than for SFT and YG because of a greater proportion of the variance explained by the study effect. The proportion of model variance explained by LMA (Model 4) was very small (5%) confirming results by *Powell et al.* [10] who showed that LMA accounted for less than 6% of the variation in chemical carcass composition (protein, fat) when animals of widely varying weights were included. There was no significant relationship between ConS and adipose tissue weight contrary to FS (Model 6) but the proportion of variance explained by FS was 17% which is very low compared with other indicators (YG, SFT, Mar). These results were similar to those of *Indurain et al.* [11] who showed that fatness

score was more efficient than conformation for predicting carcass composition.

The first driver of changes in carcass muscle weight was HCW (Model 7), explaining a high proportion of the variance (from 64 to 89 %) in all models (Models 8 to 13), while indicators contributed for only 0 to 21%. In general, the study effect was lower in muscle relationships (Models 7 to 13) than in adipose tissue relationships (Models 1 to 6) indicating that carcass muscle weight was less affected by other variation factors (such as feeding conditions) than adipose tissue [3]. Again, the YG and SFT were the most highly related to changes in carcass muscle weight (Models 8, 9), with the highest adjusted R^2 and among the lowest RMSE and MSEP. By contrast, Mar and LMA were less related to changes of muscle weight (Models 10, 11). There was a significant relationship between ConS and muscle weight (Model 12) but it was associated with a high error of prediction (29% of muscle weight). The relationships between muscle and FS (Model 13) was better than that with ConS. but these two scores (ConS and FS) were poorly related to changes in carcass muscle weights, as already noted for adipose tissues.

IV. CONCLUSION

Our results adjusted on finished cattle data showed that indirect measurements on carcass particularly fat thickness and yield grade can be good indicators of changes in tissue composition of the carcass. The exploration of these relationships should be completed by taking into account other factors of variation (breed, sex...).

REFERENCES

1. Powell, W. E. & Huffman, D. L. (1968). An evaluation of quantitative estimates of beef carcass composition. *Journal Animal Science* 27: 1554-1558.
2. Lunt, D. K., Smith, G. C., McKeith, F. R., Savell, J. W., Riewe, M. E., Horn, F. P. & Coleman, S. W. (1985). Techniques for Predicting Beef Carcass Composition. *Journal Animal Science* 60: 1201-1207.
3. Robelin, J. (1986). Composition corporelle des bovins: évolution au cours du développement et différences entre races. Thèse de Doctorat d'Etat, Université Clermont II, no. E 368, 392 pp.
4. Al-Jammas, M., Vernet, J., Agabriel, J. & Ortigues-Marty, I. (2014). Comparaison de Methodes d'évaluation de la composition Chimique de la carcasse chez les bovins par Méta-analyse. In *Viandes Produits Carnés; Hors Série*, 115-116.
5. De Boer, H., Dumont, B. L., Pomeroy, R. W. & Weniger, J. H. (1974). Manual of E.A.P.P reference methods for the assessment of carcass characteristics in Cattle. *Livestock Production Science* 1: 151-164.
6. Sauvant, D., Schmidely, P., Daudin, J. J. & St-Pierre, N. R. (2008). Meta-analyses of experimental data in animal nutrition. *Journal Animal Science* 2: 1203-1214.
7. Kobayashi, K. & Salam, M. U. (2000). Comparing Simulated and Measured Values Using Mean Squared Deviation and its Components. *Agronomy Journal* 92: 345-352.
8. Robelin, J. & Casteilla, L. (1990). Différenciation, croissance et développement du tissu adipeux. *INRA production animales* 3(4) : 243-252.
9. O'Mara, F. M., Williams, S. E., Tatum, J. D., Hilton, G. G., Pringle, T. D., Wise, J. W. & Williams, F. L. (1998). Prediction of Slaughter Cow Composition Using Live Animal and Carcass Traits. *Journal Animal Science* 76: 1594-1603.
10. Powell, W. E. & Huffman, D. L. (1973). Prediction chemical composition of beef carcasses from easily obtainable carcass variables. *Journal Animal Science* 36: 1069-1076.
11. Indurain, G., Carr, T. R., Goni, M. V., Insausti, k. & Beriain, M. J. (2009). The relationship of carcass measurements to carcass composition and intramuscular fat in Spanish beef. *Meat Science* 82:155-161.