



Efficacy of *Streptomyces* spp. strains against different strains of *Botrytis cinerea*

Sawai Boukaew, Poonsuk Prasertsan, Claire Troulet, Marc Bardin

► To cite this version:

Sawai Boukaew, Poonsuk Prasertsan, Claire Troulet, Marc Bardin. Efficacy of *Streptomyces* spp. strains against different strains of *Botrytis cinerea*. 13. IOBC-WPRS Meeting of the working group "Biological control of fungal and bacterial plant pathogens". Biocontrol of plant diseases: "from the field to the laboratory and back again", Jun 2014, Uppsala, Sweden. 2014. hal-02740549

HAL Id: hal-02740549

<https://hal.inrae.fr/hal-02740549>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Efficacy of *Streptomyces* spp. strains against different strains of *Botrytis cinerea*

Sawai Boukaew^{1,2}, Poonsuk Prasertsan¹, Claire Troulet², Marc Bardin², Department of Industrial Biotechnology, Faculty of Agro-Industry, Prince of Songkla University, Songkhla 90112, Thailand,
²INRA, UR407, Plant Pathology Unit, Domaine St Maurice, F-84140 Montfavet, France. E-mail: marc.bardin@avignon.inra.fr

Grey mould caused by the fungus *Botrytis cinerea* is an economically important disease in numerous crops. Biocontrol is a promising method to control the disease. Species of *Streptomyces* are potential biological control agents since they are ubiquitous in the environment and many of them produce various secondary metabolites with diverse biological activities including the ability to inhibit this plant pathogenic fungus. Strains RM-1-138 and RL-1-178 of *S. philanthi* and SS-2-243 of *S. mycarofaciens*, isolated from the rhizosphere soil of chili peppers grown in southern Thailand, have shown a good efficacy to control *Sclerotium rolfsii*, *Ralstonia solanacearum* and *Rhizoctonia solani* but their effect against *B. cinerea* is not known.

In this study we evaluated the efficacy of the three strains of *Streptomyces* spp. against *B. cinerea* *in vitro* and on tomato plants. Results indicated that the three strains inhibit the growth of *B. cinerea* in Petri plates and have a significant protective efficacy, although variable between strains of *Streptomyces* spp., against *B. cinerea* on tomato plant.

To assess the possible variability in susceptibility to these antagonistic strains in populations of *B. cinerea*, the protective efficacy of these bacteria were evaluated against 41 strains differing in their geographic origin, host of isolation and level of aggressiveness. Results based on confrontation tests in Petri plates suggest a diversity in the sensitivity of the different strains of *B. cinerea* to these biological control agents. The relevance of these results to ensure durability of the efficacy of these biological control agents will be discussed.



International Organisation for Biological and Integrated Control of Noxious Animals and Plants (IOBC)

West Palaearctic Regional Section (WPRS)

Organisation Internationale de Lutte Biologique et Intégrée contre les Animaux et les Plantes Nuisibles (OILB)

Section Régionale Ouest Paléarctique (SRPOP)

XIII Meeting of the Working Group Biological control of fungal and bacterial plant pathogens

Viticulture

Horticulture

Agriculture

Forestry

Biocontrol of Plant Diseases: “From the field to the laboratory and back again”



Uppsala, Sweden
SLU, Swedish University of
Agricultural Sciences



15 – 18 June 2014

Programme Abstracts Participants

