



HAL
open science

Response of black poplar (*Populus nigra* L.) to hydrogeomorphological constraints: a semi-controlled ex situ experiment

Virginia Garófano-Gómez, Dov Corenblit, Johannes Steiger, Bruno B. Moulia, Stéphane S. Ploquin, Patrice Chaleil, Olivier Forestier, André Evette, Eduardo Gonzalez, Borbála Hortobágyi, et al.

► To cite this version:

Virginia Garófano-Gómez, Dov Corenblit, Johannes Steiger, Bruno B. Moulia, Stéphane S. Ploquin, et al.. Response of black poplar (*Populus nigra* L.) to hydrogeomorphological constraints: a semi-controlled ex situ experiment. *I.S. Rivers*, 2e conférence internationale "recherches et actions au service des fleuves et grandes rivières", Jun 2015, Lyon, France. La ZABR et le GRAIE, 324 p., 2015, Recherches et actions au service des fleuves et grandes rivières. hal-02740599

HAL Id: hal-02740599

<https://hal.inrae.fr/hal-02740599>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RESPONSE OF BLACK POPLAR (*POPULUS NIGRA* L.) TO HYDROGEOMORPHOLOGICAL CONSTRAINTS: A SEMI-CONTROLLED *EX SITU* EXPERIMENT

Réponse du peuplier noir (*Populus nigra* L.) aux contraintes hydro-géomorphologiques : une expérimentation *ex situ* semi-contrôlée


1 Evolutionary feedback between woody riparian species and hydrogeomorphological constraints

- Hydrogeomorphological factors (topography, flow and sediment transport regimes) control vegetation dynamics in riparian ecosystems → but vegetation also has an impact on these factors, which in turn causes an effect on the plant phenotype.
- Concepts:** 'ecosystems engineers' and 'positive niche construction'.
- At an evolutionary timescale, this **reciprocal interaction** has promoted the selection of certain **plant traits** to increase the persistence of woody riparian species within fluvial environments.



2 The biogeomorphological life cycle of black poplar (*Populus nigra* L.)

- Black poplar is a keystone ecosystem engineer species. Specific ranges of hydrogeomorphological conditions control the successive phases of its entire life cycle.
- Hypothesis:** the impact of poplars on the landform structure modulates its own growth performance, biomass and architecture until it reaches sexual maturity.





3 Semi-controlled *ex situ* experiment

- 3.1 Objective:** To quantify **key response functional traits** (morphological and biomechanical) of *Populus nigra* L. cuttings to simulated hydrogeomorphological constraints (**drag force** and **sediment burial**) as well as to dissociate the specific responses to them.
- 3.2 Experimental design:** 128 stem cuttings of *P. nigra* (variety Jean Pourtet) were measured, planted in permeable bags with an irrigation system attached and randomly assigned to one of the 4 treatments.



3.2.1 Treatments:



- The weight (T.1) will be modulated imitating the shape of an average hydrograph of a Spring flood in the Garonne River (where the clone Jean Pourtet comes from).
- The burial (T.2) will be applied during the recession limb of the curve.

- Temporal sequence** of expected above-ground and below-ground plant development according to the application of treatments. (Experimentation from March to Sept. 2015)




3.2.2 Morphological and biomechanical traits:

| Above-ground traits | Below-ground traits | Ratios |
|---|--|--|
| <ul style="list-style-type: none"> Number of shoots Max. plant height Root collar diameter Diameter at middle mature height Tapering Inclination of the main stem Average leaf area Specific leaf area Above-ground dry mass Frontal surface area Pulling force* Flexibility* | <ul style="list-style-type: none"> Initial diameter (cutting) Initial weight (cutting) N° first order roots N° structural roots N° basal, lateral and superficial roots Root diameter Insertion angle Root length by diameter class Max. and mean root length Below-ground dry mass N° 'shear' and 'broken' roots* Diameter of 'shear' and 'broken' roots* | <ul style="list-style-type: none"> Root mass fraction Shoot mass fraction Elongation ratio Shoot to root ratio Fine/structural roots Leaf area to root length ratio Root weight/n° of tips Roots extracted/remaining in the bag* |


* Traits from *winching test*.

3.2.3 Preliminary results: First partial harvest

- 12 plants** were destructively sampled to test the methodology of extraction, conservation and sub-sampling.



- The **growth is optimum** but some differences are evident depending on the original size of the cutting and the mother tree they come from.



- Analysis of root length confirms that roots could have different **functions**: anchorage (basal roots) and absorption (lateral roots).

The quantification of functional response traits of *P. nigra* will enhance our understanding of fundamental biogeomorphic interactions and its implication for the restoration of river systems.

Acknowledgements: to Catherine Cochard, Olivier Voldoire, Adrià Masip, Mohamed Abadi, Christophe Corona, Jérôme Lopez, Valerie Legué, Frédéric Julien and Marc Villar for their support during this experimentation.

Virginia Garófano-Gómez^{1,2}; Dov Corenblit^{1,2}; Johannes Steiger^{1,2}; Bruno Mouliat³; Stéphane Ploquin³; Patrice Chaleil³; Olivier Forestier⁴; André Evette⁵; Eduardo González^{6,7}; Borbála Hortobágyi^{1,2} and Luc Lambs⁶

¹Université Clermont Auvergne, UBP, MSH, Clermont-Ferrand, France (corresponding author: virginia.garofano_gomez@univ-bpclermont.fr). ²CNRS, UMR 6042, GEOLAB – Laboratoire de géographie physique et environnementale, Clermont-Ferrand, France. ³INRA Clermont Ferrand, UMR547 Laboratoire de physique et physiologie intégrative de l'arbre fruitier et forestier, Site de Crouël, Clermont Ferrand, France. ⁴Pépinière Forestière de l'Etat, DRAAF Pays-de-la-Loire, Guémené Penfao, France. ⁵Irstea, UR EMGR, Saint-Martin-d'Hères, France. ⁶Ecolab CNRS, Toulouse, France. ⁷Department of Biological Sciences, University of Denver, Canada.