



**HAL**  
open science

## Vulnerability to embolism in herbaceous and secondarily woody species

Frederic Lens, Hervé H. Cochard, Catherine Picon-Cochard, Steven Jansen, Thibaud Chauvin, Larissa Chacon Doria, Marcelino del Arco, Sylvain S. Delzon

### ► To cite this version:

Frederic Lens, Hervé H. Cochard, Catherine Picon-Cochard, Steven Jansen, Thibaud Chauvin, et al.. Vulnerability to embolism in herbaceous and secondarily woody species. Xylem International Meeting, Sep 2015, Bordeaux, France. 86 p. hal-02740600

**HAL Id: hal-02740600**

**<https://hal.inrae.fr/hal-02740600>**

Submitted on 2 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Vulnerability to embolism in herbaceous and secondarily woody species

Frederic Lens<sup>1\*</sup>, Hervé Cochard<sup>2</sup>, Catherine Picon-Cochard<sup>3</sup>, Steven Jansen<sup>4</sup>, Thibaud Chauvin<sup>1,5</sup>, Larissa Chacon Doria<sup>1</sup>, Marcelino del Arco<sup>6</sup> and Sylvain Delzon<sup>5</sup>.

<sup>1</sup> Naturalis Biodiversity Center, Leiden University, P.O. Box 9517, 2300RA Leiden, the Netherlands

<sup>2</sup> INRA, UMR 547 PIAF, F-63100 Clermont-Ferrand, France.

<sup>3</sup> INRA UR874 Grassland Ecosystem Research F-63039 Clermont-Ferrand cedex 2, France.

<sup>4</sup> Institute for Systematic Botany and Ecology, Ulm University, Albert-Einstein-Allee 11, D-89081, Ulm, Germany.

<sup>5</sup> INRA, University of Bordeaux, UMR BIOGECO, F-33450 Talence, France.

<sup>6</sup> Department of Plant Biology (Botany), La Laguna University, 38071 La Laguna, Tenerife, Spain.

\*Author for correspondence: frederic.lens@naturalis.nl

### Abstract:

There is a common idea that most herbaceous species refill their embolized vessels each night via positive root pressure, meaning that there is no need to become resistant to embolism. Our study assesses the range in vulnerability to embolism in herbaceous species, and evaluates whether closely related secondarily woody species are always more embolism resistant than their herbaceous ancestors. Therefore, three non-related plant groups are investigated: 1) grass species, 2) the Gentianaceae sister pair *Blackstonia* (herbaceous) and *Ixanthus* (woody), and 3) the daisy lineage (Asteraceae) with herbaceous (*Leucanthemum*, *Chamaemelum*) and woody members (*Argyranthemum*). Surprisingly, the grass dataset shows an extraordinary range in P50, ranging from 0.5 – 7.5MPa. This suggests that at least in the resistant species daily refilling does not occur, although this needs to be quantified. The anatomy of the grass stems shows a tight trade-off between P50 and stem lignification. Within the daisy lineage, all the woody species are more embolism resistant than their herbaceous relatives, but the *Blackstonia-Ixanthus* sister group has a similar P50 value (-4.5MPa), showing that more woodiness/lignification does not always go hand in hand with more embolism resistance.