



Prospective associations between vitamin D status, vitamin D-related gene polymorphisms, and risk of tobacco-related cancers

Melanie Deschasaux, Paule Latino Martel, Jean-Claude Souberbielle, Angela Sutton, Nathalie Charnaux, Nathalie Druesne Pecollo, Pilar Galan, Serge Hercberg, Sigrid Le Clerc, Emmanuelle Kesse-Guyot, et al.

► To cite this version:

Melanie Deschasaux, Paule Latino Martel, Jean-Claude Souberbielle, Angela Sutton, Nathalie Charnaux, et al.. Prospective associations between vitamin D status, vitamin D-related gene polymorphisms, and risk of tobacco-related cancers. IARC's 50th Anniversary, Jun 2016, Lyon, France. , 518 p., 2016, IARC's 50th Anniversary "Cancer: Occurrence, Causes, Prevention". Abstracts. hal-02740715

HAL Id: hal-02740715

<https://hal.inrae.fr/hal-02740715>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

B-068 - Prospective Associations Between Vitamin D Status, Vitamin D-Related Gene Polymorphisms, And Risk Of Tobacco-Related Cancers

MÉLANIE DESCHASAUX, SORBONNE PARIS CITÉ EPIDEMIOLOGY AND STATISTICS RESEARCH CENTER (CRESS), INSERM U1153, INRA U1125, CNAM, PARIS 13 UNIVERSITY, FRANCE

LATINO-MARTEL P.¹, SOUBERBIELLE J.², SUTTON A.³, CHARNAUX N.³, DRUESNE-PECOLLO N.¹, GALAN P.¹, HERCBERG S.^{1,4}, LE CLERC S.⁵, KESSE-GUYOT E.¹, EZZEDINE K.⁶, TOUVIER M.¹

¹ Sorbonne Paris Cité Epidemiology and Statistics Research Center (CRESS), Inserm U1153, Inra U1125, Cnam, Paris 13 University, Nutritional Epidemiology Research Team (EREN), 93017 Bobigny, France

² Physiology Department, Necker Hospital, Inserm U845, 75015 Paris, France

³ Biochemistry Department, Jean Verdier Hospital, Inserm U698, Paris 13 University, 93140 Bondy, France

⁴ Public Health Department, Avicenne Hospital, 93017 Bobigny, France

⁵ Conservatoire National des Arts et Métiers (CNAM), Genomics, Bioinformatics and Applications Team (EA4627), 75003 Paris, France

⁶ Dermatology Department, Saint André Hospital, 33000 Bordeaux, France

Purpose:

Experimental evidence has suggested that vitamin D may be protective against tobacco-related cancers through the inhibition of the formation of tumors induced by tobacco carcinogens. To our knowledge, only one previous epidemiologic study investigated the association between vitamin D status and tobacco-related cancer risk, and no study has focused on vitamin D-related gene polymorphisms. Our objective was to prospectively study the association between plasma 25-hydroxyvitamin D [25(OH)D] concentrations, vitamin D-related gene polymorphisms (VDR, CYP24A1, GC, RXR, CaSR), and risk of tobacco-related cancers.

Methods:

A total of 209 tobacco-related cancers were diagnosed within the SU.VI.MAX (Supplémentation en vitamines et minéraux antioxydants) cohort (1994–2007) and were matched with 418 controls as part of a nested case-control study. Tobacco-related cancers (i.e., cancers for which tobacco is one of the risk factors) included several sites in the respiratory, digestive, reproductive, and urinary systems. Plasma total 25(OH)D concentration and selected gene polymorphisms were assessed on samples obtained at baseline. Conditional logistic regression models were computed.

Results:

A 25(OH)D concentration ≥ 30 ng/mL was associated with reduced risk of tobacco-related cancers ($OR_{\geq 30 \text{ vs. } < 30 \text{ ng/mL}} = 0.59$ (0.35–0.99); $P = 0.046$). This association was observed in former and current smokers ($OR_{\geq 30 \text{ vs. } < 30 \text{ ng/mL}} = 0.43$ (0.23–0.84); $P = 0.01$) but not in never smokers ($P = 0.8$). The vitamin D receptor (VDR) FokI AA genotype and retinoid X receptor (RXR) rs7861779 TT genotype were associated with increased risk of tobacco-related cancers.

Conclusions:

In this prospective study, high vitamin D status [25(OH)D concentration ≥ 30 ng/mL] was associated with decreased risk of tobacco-related cancers, especially in smokers. These results, which are supported by mechanistic plausibility, suggest that vitamin D may contribute to the prevention of tobacco-induced cancers in smokers and deserve additional investigation.

Am J Clin Nutr 2015, 102:1207–15.

Funding source:

French Research Institute for Public Health (IRESP), grant number AAR201206

Cancéropôle Ile-de-France, Ile-de-France Region, PhD grant (Mélanie Deschasaux)