

Transfer equations for modeling interrill erosion

Amina Nouhou Bako, Frédéric Darboux, François James, Carine Lucas

▶ To cite this version:

Amina Nouhou Bako, Frédéric Darboux, Francois James, Carine Lucas. Transfer equations for modeling interrill erosion. EGU, European Geosciences Union General Assembly 2016, Apr 2016, Vienne, Austria. hal-02740915

HAL Id: hal-02740915 https://hal.inrae.fr/hal-02740915v1

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Geophysical Research Abstracts Vol. 18, EGU2016-494, 2016 EGU General Assembly 2016 © Author(s) 2015. CC Attribution 3.0 License.



Transfer equations for modeling interrill erosion

Nouhou Bako Amina (1,2), Darboux Frédéric (1), James François (2), and Lucas Carine (2) (1) Institut National de la Recherche Agronomique, Science du Sol, orléans, France (amina.nouhou-bako@orleans.inra.fr), (2) Université d'Orléans, MAPMO, Orléans, France

Numerous models are available for matter transfer along an hillslope. They are usually process-specific, requiring to use several models to simulate transfers along an hillslope. To overcome this issue, we develop a new model valid for chemical (nutrients, pollutants, dissolved carbon) and particle transfers by water. It is able to simulate both interrill and rill erosion. This new equation encompasses the previous models of Gao et al. (2004), Hairsine and Rose (1992, 1991) and Lajeunesse et al. (2013) in a single and unified form. We show that it can account for multi-class particle transport able to simulate both linear and non-linear behaviors. Surface conditions (crusts) is accounted for, making possible for space and time changes of soil properties.

For the calibration of the model, specific laboratory experiments have been carried out to validate the effect of rainfall on travel distance of particles. These experiments allow to separate detachment by raindrops from the agitation of the flow by the drops. Different particle sizes and rainfall kinetic energies are investigated. The results assess the exact role of rainfall on sediment transport. Our new model is able to represent adequately these experimental results.