



HAL
open science

Tolerance of maize to drought: a matter of aquaporins?

Linda Jeanguenin, Llorenç Cabrera Bosquet, Claude Welcker, Francois F. Tardieu, François Chaumont

► To cite this version:

Linda Jeanguenin, Llorenç Cabrera Bosquet, Claude Welcker, Francois F. Tardieu, François Chaumont. Tolerance of maize to drought: a matter of aquaporins?. Recent progress in drought tolerance from genetics to modelling, Jun 2015, Montpellier, France. 2015. hal-02740989

HAL Id: hal-02740989

<https://hal.inrae.fr/hal-02740989>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Interactive exhibit

Tolerance of maize to drought: a matter of aquaporins?

Linda Jeanguenin ^{1,a}, Llorenç Cabrera-Bosquet ^{2,b}, Claude Welcker ^{3,b}, François Tardieu ^{4,b}, François Chaumont ^{5,a}

^a Institut des Sciences de la Vie, Université catholique de Louvain, Croix du sud
4-L7.07.14, 1348 Louvain-la-Neuve, Belgium

^b INRA-LEPSE, 2 place Viala, 34060 Montpellier, France

linda.jeanguenin@uclouvain.be

Plant aquaporins play key roles in the cellular water permeability by enhancing the transport of water molecules across lipophilic membranes. In maize, four aquaporin subfamilies were identified, among them were found the PIPs standing for plasma membrane intrinsic proteins. Despite the fact that PIP aquaporins are water channels, their function in the plant response to water deficit is not well characterized (Chaumont and Tyerman, 2014). Gene candidate approaches based on the overexpression of one specific *PIP* gene can lead to either deleterious or beneficial effects for plants under water shortage. Microarray global expression studies on five *Arabidopsis thaliana* accessions showed a decreased transcription of most *PIP* genes under progressive drought (Alexandersson et al., 2010). In addition, physiological and pharmacological approaches conducted among several crops showed that aquaporins are involved in both leaf elongation and transpiration rates in response to atmospheric water deficit (Sadok and Sinclair, 2010; Caldeira et al., 2014). Altogether these data indicate that aquaporins are involved in the plant response to water limitation. Under the frame of the European DROPS project, the expression of 12 *PIP* isoforms was investigated by quantitative PCR in 30 maize lines grown under two contrasted water regimes in the PhenoArch phenotyping platform (LEPSE, Montpellier) where biomass, water use efficiency and transpiration were measured in parallel. Under water limitation, the mRNA level of *PIPs* belonging to the PIP1 group was weakly regulated while it was down-regulated for most *PIP2s*, in both leaf elongation and mature zones. *PIP* expression pattern to water deficit also relied upon the plant life cycle. Furthermore, principal component analysis revealed that expression of most *PIPs* contributed to the first component while biomass and hydraulic parameters contributed to the second component in well-watered conditions. Under water deficit, this distribution changed indicating that aquaporin expression adjustment to water limitation is involved in the regulation of these physiological parameters.

Keywords: Aquaporins, Leaf, Maize (*Zea mays*), Plasma membrane intrinsic protein (PIP), Water deficit