



HAL
open science

Soil moisture estimation across scales with mobile sensors for cosmic-ray neutrons from the ground and air

Matthias Cuntz, Martin Schrön

► To cite this version:

Matthias Cuntz, Martin Schrön. Soil moisture estimation across scales with mobile sensors for cosmic-ray neutrons from the ground and air. EGU, European Geosciences Union General Assembly 2016, European Geosciences Union (EGU). AUT., Apr 2016, Vienne, Austria. 1 p. hal-02741481

HAL Id: hal-02741481

<https://hal.inrae.fr/hal-02741481>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Soil Moisture Estimation Across Scales with Mobile Sensors for Cosmic-Ray Neutrons from the Ground and Air

Martin Schrön (1), Mandy Köhler (1), Lutz Bannehr (2), Markus Köhli (3), Benjamin Fersch (4), Corinna Rebmann (1), Juliane Mai (1), Matthias Cuntz (1), Simon Kögler (1), Ingmar Schröter (1), Ute Wollschläger (1), Sascha Oswald (5), Peter Dietrich (1), and Steffen Zacharias (1)

(1) UFZ - Helmholtz Centre for Environmental Research, Computational Hydrosystems, Leipzig, Germany (martin.schroen@ufz.de), (3) Physics Institute, University of Heidelberg, (2) Hochschule Anhalt, Institut für Geoinformation und Vermessung, Dessau, (4) KIT - IMK-IFU Institute of Meteorology and Climate Research, (5) Institut für Erd- und Umweltwissenschaften, University of Potsdam

Soil moisture is a key variable for environmental sciences, but its determination at various scales and depths is still an open challenge.

Cosmic-ray neutron sensing has become a well accepted and unique method to monitor an effective soil water content, covering tens of hectares in area and tens of centimeters in depth. The technology is famous for its low maintenance, non-invasiveness, continuous measurement, and most importantly its large footprint and penetration depth. Being more representative than point data, and finer resolved plus deeper penetrating than remote-sensing products, cosmic-ray neutron derived soil moisture products provide unrivaled advantage for agriculture, regional hydrologic and land surface models.

The method takes advantage of omnipresent neutrons which are extraordinarily sensitive to hydrogen in soil, plants, snow and air. Unwanted hydrogen sources in the footprint can be excluded by local calibration to extract the pure soil water information. However, this procedure is not feasible for mobile measurements, where neutron detectors are mounted on a car to do catchment-scale surveys.

As a solution to that problem, we suggest strategies to correct spatial neutron data with the help of available spatial data of soil type, land use and vegetation. We further present results of mobile rover campaigns at various scales and conditions, covering small sites from 0.2 km² to catchments of 100 km² area, and complex terrain from agricultural fields, urban areas, forests, to snowy alpine sites.

As the rover is limited to accessible roads, we further investigated the applicability of airborne measurements. First tests with a gyrocopter at 150 to 200m heights proofed the concept of airborne neutron detection for environmental sciences. Moreover, neutron transport simulations confirm an improved areal coverage during these campaigns.

Mobile neutron measurements at the ground or air are a promising tool for the detection of water sources across many scales. The method has a great potential to improve spatial performance of hydrological models, and help to assess regional soil moisture states for agriculture and flood risk management.