

Carbon and water cycling in flooded and rainfed rice (Oryza Sativa) ecosystem: Disentangling agronomical and ecological aspects of water use efficiency

Bhone Nay-Htoon, Xue Wei, Dubbert Maren, Steve Lindner, Matthias Cuntz, Ko Jonghan, John Tenhunen, Christiane Werner

► To cite this version:

Bhone Nay-Htoon, Xue Wei, Dubbert Maren, Steve Lindner, Matthias Cuntz, et al.. Carbon and water cycling in flooded and rainfed rice (Oryza Sativa) ecosystem: Disentangling agronomical and ecological aspects of water use efficiency. EGU 2015, European Geophysical Union General Assembly 2015, Apr 2015, Vienne, Austria. 2015. hal-02741490

HAL Id: hal-02741490 https://hal.inrae.fr/hal-02741490v1

Submitted on 3 Jun 2020 $\,$

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Carbon and water cycling in flooded and rainfed rice (Oryza Sativa) ecosystem: Disentangling agronomical and ecological aspects of water use efficiency

Bhone Nay-Htoon (1), Wei Xue (2), Maren Dubbert (1), Steve Lindner (2), Matthias Cuntz (3), Jonghan Ko (4), John Tenhunen (2), and Christiane Werner (1)

(1) Department of Agroecosystem Research, BayCEER, University of Bayreuth, Bayreuth, Germany, (2) Department of Plant Ecology, BayCEER, University of Bayreuth, Bayreuth, Germany, (3) Computational Hydrosystems, Helmholtz Centre for Environmental Research (UFZ), Leipzig, Germany, (4) Division of Applied Plant Science, Chonnam National University, Gwangju, Republic of Korea

Agricultural crops play an important role in the global carbon and water cycling process and there is intense research to understand and predict carbon and water fluxes, productivity and water use of cultivated crops under climate change. Mechanistic understanding of the trade of between ecosystem water use efficiency and agronomic water use efficiency to maintain higher crop yield and productive water loss is necessary for the ecosystem sustainability. We compared water and carbon fluxes of paddy and rainfed rice by canopy scale gas exchange measurements, crop growth, and daily evapotranspiration, transpiration and carbon flux modeling.

According to our findings, evaporation contributed strongly (maximum 100% to minimum 45%) to paddy rice evapotranspiration while transpiration of rainfed is almost 50 % of daily evapotranspiration. Water use efficiency (WUE) was higher in rainfed rice both from an agronomic (WUEagro, i.e. grain yield per evapotranspiration) and ecosystem (WUEeco, i.e. gross primary production per evapotranspiration) perspective. However, rainfed rice showed also high ecosystem respiration losses and a slightly lower crop yield, demonstrating that higher WUE in rainfed rice comes at the expense of higher respiration losses of assimilated carbon and lower plant production, compared to paddy rice. Our results highlighted the need to partition water and carbon fluxes to improve our mechanistic understanding of water use efficiency and environmental impact of different agricultural practices.

Keywords: Rainfed rice, Paddy rice, water use efficiency, Transpiration/Evapotranspiration, ecosystem WUE, agronomic WUE, Evapotranspiration