



HAL
open science

Crop yields, soil organic carbon and soil nitrogen content change under climate change

B. Dumont, B. Basso, I. Shcherbak, S. Asseng, Simona Bassu, K. Boote, D. Cammarano, Giovanni de Sanctis, Jean-Louis Durand, F. Ewert, et al.

► **To cite this version:**

B. Dumont, B. Basso, I. Shcherbak, S. Asseng, Simona Bassu, et al.. Crop yields, soil organic carbon and soil nitrogen content change under climate change. iCROP 2016 International Crop Modelling Symposium, Leibniz-Zentrum für Agrarlandschaftsforschung = Leibniz Centre for Agricultural Landscape Research, Leibniz Association (ZALF). DEU., Mar 2016, Berlin, Germany. 437 p. <hal-02741515>

HAL Id: hal-02741515

<https://hal.inrae.fr/hal-02741515v1>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Crop yields, soil organic carbon and soil nitrogen content change under climate change

B. Dumont^{1,*} - *B. Basso*¹ - *I. Shcherbak*¹ - *S. Asseng*² - *S. Bassu*³ - *K. Boote*⁴ -
*D. Cammarano*⁵ - *G. de Sanctis*⁶ - *J.-L. Durand*⁷ - *F. Ewert*⁸ - *S. Gayler*⁹ - *P. Grace*¹⁰ -
*R. Grant*¹¹ - *J. Kent*¹² - *P. Martre*¹³ - *C. Nendel*¹⁴ - *K. Paustian*² - *E. Priesack*¹⁵ -
*D. Ripoche*⁶ - *A. Ruane*¹⁷ - *P. Thorburn*¹⁸ - *J. Hatfield*¹⁹ - *J. Jones*²⁰ - *C. Rosenzweig*¹⁷

¹ Department of Geological Sciences, Michigan State University, MI, USA - ² Agricultural and Biological Engineering Department, University of Florida, Gainesville, FL, USA - ³ INRA-AgroParisTech, Thiverval-Grignon, France - ⁴ Department of Agronomy, University of Florida, Gainesville, FL, USA - ⁵ The James Hutton Institute, Invergowrie, Scotland, UK - ⁶ European Commission - Joint Research center, Ispra, Italy - ⁷ INRA-URP3F, Lusignan, France - ⁸ Institute of Crop Science and Resource Conservation (INRES), University of Bonn, Bonn, Germany - ⁹ WESS-Water and Earth System Science Competence Cluster, University of Tübingen, Tübingen, Germany - ¹⁰ Institute for Future Environments, Queensland University of Technology, Brisbane, Queensland, Australia - ¹¹ Institute of Crop Science and Resource Conservation (INRES), University of Bonn, Bonn, Germany - ¹² Natural Resource Ecology Laboratory, Colorado State University, Fort Collins, CO, USA - ¹³ INRA, UMR759 LEPSE, Montpellier, France - ¹⁴ Institute of Landscape Systems Analysis, ZALF, Leibniz-Centre for Agricultural Landscape Research, Muencheberg, Germany - ¹⁵ Institute of Soil Ecology, Helmholtz Zentrum München—German Research Center for Environmental Health, Neuherberg, Germany - ¹⁶ INRA-AGROCLIM, Avignon, France - ¹⁷ Climate Impacts Group, NASA Goddard Institute for Space Studies, New York, NY, USA - ¹⁸ CSIRO Ecosystem Sciences, Dutton Park, Queensland, Australia - ¹⁹ USDA-ARS National Soil Tilth Laboratory for Agriculture and the Environment, Ames, IA, USA - ²⁰ Department of Agricultural and Biological Engineering, University of Florida, Gainesville, FL, USA.

* 288 Farm Lane, Dpt. Geological Sciences, Natural Sciences Bldg., # 306, Michigan State University, MI, USA
Tel : +1-517-353-3472 – E-mail : bdumont@msu.edu

Introduction

It has been demonstrated that model ensembles is an efficient way to reduce the uncertainty associated with climate change impact on crop growth. (Asseng et al., 2015). Using such this approach, wheat and maize grain yields response to temperature increase were simulated by Asseng et al., 2015 and Bassu et al., 2014 using annually re-initialized soil conditions (soil water and nitrogen). Bassu et al., 2015, showed that by running models in a continuous mode, yield results differed from the annual reinitialized runs. In this study, we present the results of continuous model run of the AgMIP wheat- and maize-pilot under temperature and CO₂ changes and different management practices.

Materials and Methods

Five maize models and seven wheat models were run using a continuous mode under conventional tillage and no-till using the same factorials temperature, CO₂, rain and nitrogen fertilization levels of the AgMIP pilots (Asseng et al., 2015 and Bassu et al., 2014). We evaluate the range of response provided by the different models soil organic carbon (SOC) dynamic, soil nitrogen (N-NO₃⁻) and water dynamics under the maize-fallow and wheat-fallow crop rotations.

Results and Discussion

Under continuous running mode, models agreed in the direction of the changes for soil N-NO₃⁻ and SOC under the different temperature treatments. Furthermore, soil N-NO₃⁻ was found to increase and SOC was found to decrease with temperature increases. Consequently, important differences on simulated yields were observed between reinitialized and continuous run of the model. Fig. 1 compares the average model results under different CO₂ treatments and soil management practices for the continuous run. Whatever the temperature, yields were overall higher when CO₂ increased and lower when practicing no-till. Soil N-NO₃⁻ was lower under higher CO₂ concentration and was found to increase with temperature.




Figure 1. Wheat yield (A-B) and soil N-NO₃⁻ (C-D) content under different temperature (-3 to +9°C) treatments. Impact of CO₂ concentration (A-C) and tillage practice (B-D).

Conclusions

Continuous running mode of crop models appeared as a promising way to better understand the interactions between soil, climate and crop management. Such an approach is a promising mean to conceive crop and soil management strategies able to mitigate adverse climate change impacts.

References

- Asseng, S., Ewert, F., Martre, P. et al., (2015). Nature Clim. Change 5: 143-147.
 Bassu, S., Brisson, N., Durand, J.L., Boote, K. et al., (2014). Global Change Biology 20: 2301-2320.
 Basso B. et al., (2015). Plos One 10(6): e0127333.