



**HAL**  
open science

## Drought influence on carbon and water cycling in a Mediterranean *Quercus suber* L. woodland during the drought year 2012

Arndt Piayda, Maren Dubbert, Corinna Rebmann, Olaf Kolle, Filipe Costa E Silva, Alexandra Correia, Joao Santos Pereira, Christiane Werner, Matthias Cuntz

### ► To cite this version:

Arndt Piayda, Maren Dubbert, Corinna Rebmann, Olaf Kolle, Filipe Costa E Silva, et al.. Drought influence on carbon and water cycling in a Mediterranean *Quercus suber* L. woodland during the drought year 2012. EGU 2014, European Geosciences Union General Assembly 2014, Apr 2014, Vienne, Austria. 2014. hal-02741526

**HAL Id: hal-02741526**

**<https://hal.inrae.fr/hal-02741526>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



## Drought influence on carbon and water cycling in a Mediterranean *Quercus suber* L. woodland during the drought year 2012

Arndt Piayda (1), Maren Dubbert (2), Corinna Rebmann (1), Olaf Kolle (3), Filipe Costa e Silva (4), Alexandra Correia (4), Joao Santos Pereira (4), Christiane Werner (2), and Matthias Cuntz (1)

(1) Helmholtz Centre for Environmental Research – UFZ, Department Computational Hydrosystems, Leipzig, Germany (arndt.piayda@ufz.de), (2) University of Bayreuth, Agroecosystem Research Department, Bayreuth, Germany, (3) Max Planck Institute for Biogeochemistry, Field Experiments & Instrumentation, Jena, Germany, (4) Instituto Superior de Agronomia, Department of Forestry, Technical University of Lisbon, Lisbon, Portugal

Savannah-type ecosystems account for 26-30% of the global gross primary production with water being one of the major driving factors. In Europe, savannah-type woodlands cover an area of about 2-2.5 million ha on the Iberian Peninsula. The recent past has shown there a significant decrease of precipitation in winter and spring as well as a decrease of total annual precipitation. Hence, strong effects on local water balance and carbon sink strength have been reported due to these changes in the precipitation regime.

The objective of this study is to quantify changes in the water balance, gross primary productivity and carbon sink strength of a typical Portuguese savannah-type woodland (*montado*) under the changed precipitation pattern of the drought year 2012 compared to the wet year 2011. The physiological response of the dominant tree species *Quercus suber* (L.) is evaluated, employing combined photosynthesis and stomatal conductance modelling.

Precipitation effectiveness ratio increased up to 122% in the dry year 2012 due to ground water access of the *Q. suber* trees leaving no water for ground water replenishment. By the lack of water in the upper soil and deep ground water reservoirs, the understorey and overstorey gross primary productions were reduced by 53% and 28% in 2012 compared to 2011, respectively, due to the late onset of the autumn rains in 2011 and an additional severe winter/spring drought. However, on an annual basis, the ecosystem was a carbon sink in both years, with a 61% reduced sink strength in the dry year 2012 compared to the wet 2011.

Applying a combined photosynthesis and stomatal conductance model, best model fit to gross primary productivity and transpiration of *Q. suber* trees could be achieved keeping apparent maximum carboxylation rate  $V_{c,max}$  as well as stomatal conductivity parameter  $m$  and vapor pressure deficit sensitivity parameter  $D_0$  of the stomatal conductance formulation variable. The *Q. suber* trees showed 20% reduced stomatal conductance  $g_s$  during the drought period 2012 compared to 2011 expressed as a reduction in stomatal conductance model parameter  $m$ . Hence, vapor pressure deficit sensitivity parameter  $D_0$  increased under drought conditions in order to preserve the sensitivity of  $g_s$  to vapor pressure deficit  $vpd$ . In response to reduced leaf internal  $CO_2$  availability and increased leaf temperatures, the trees strongly reduced apparent maximum carboxylation rate  $V_{c,max}$  by 39% in 2012 compared to 2011. However, stomatal response was strongest at the drought beginning while strongest reduction of  $V_{c,max}$  occurred at the end of the drought.

Our results suggest that, if the trend of decreasing annual precipitation and changed precipitation pattern on the Iberian Peninsula continues, sustained effects on local ground water reservoirs, understorey species composition and tree productivity may be expected in the long-term. To successfully model the effect of drought on the *montado* ecosystem, variable apparent maximum carboxylation rate  $V_{c,max}$ , stomatal conductivity parameter  $m$  and vapor pressure deficit sensitivity parameter  $D_0$  need to be incorporated in photosynthesis and stomatal conductance modelling.