



HAL
open science

Nickel Phytoextraction by Seven Populations of Hyperaccumulating Brassicaceae.

Marie Rue, Luciana Frota-Madeira, Jean-Louis Morel, Marie-Odile Simonnot

► **To cite this version:**

Marie Rue, Luciana Frota-Madeira, Jean-Louis Morel, Marie-Odile Simonnot. Nickel Phytoextraction by Seven Populations of Hyperaccumulating Brassicaceae.. 13th SGA Meeting “Mineral Resources in a Sustainable World, Aug 2015, Nancy, France. 2015. hal-02741676

HAL Id: hal-02741676

<https://hal.inrae.fr/hal-02741676>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License

Nickel Phytoextraction by Seven Populations of Hyperaccumulating Brassicaceae

Marie RUE^{1,2}, Luciana FROTA-MADEIRA, Jean Louis MOREL², Marie-Odile SIMONNOT¹

¹ Laboratoire Réactions et Génie des Procédés, UMR 7274 CNRS - Université de Lorraine, BP 20451, 54001 Nancy

² Laboratoire Sols et Environnement, UMR 1120 INRA - Université de Lorraine, BP 172, 54518 Vandoeuvre-lès-Nancy

marie.rue@univ-lorraine.fr

Context

Hyperaccumulator plants are able to accumulate one or several metals at high concentration. For nickel (Ni), the threshold concentration is fixed at **1000 mg kg⁻¹ of Ni** in dry matter of shoots.

Hyperaccumulator plants can be used to extract metals from contaminated soils or from soils naturally rich in Ni called ultramafic soils. Then, Ni sequestered in plants can be recovered by hydrometallurgical processes for metal recycling.

The optimisation of agronomic practices for growing these plants is a current challenge. It involves the selection of efficient plants, i.e. plants with **high biomass production** and **high concentration of metals in their shoots**.

This research is part of the LORVER project (www.lorver.org) aiming at creating a production sector of non-food plant biomass for industrial purposes, the biomass being grown on degraded sites and industrial by-products. The present objective is to evaluate the phytoextraction potential of various Ni hyperaccumulators across species, populations and individuals.

Materials and methods

ULTRAMAFIC SOIL	
pH	6.14
C/N	12.6
CEC	31.3 cmol ⁺ kg ⁻¹
[Ni] _{total}	1480 mg kg ⁻¹
[Ni] _{DTPA}	83 mg kg ⁻¹

Ultramafic soil has been sampled in the Vosges mountains. Pots were prepared with 70 g of soil; 70 plants of each population were grown during 70 d in controlled conditions.

PLANTS: 7 populations

- ✓ *Alyssum murale* (ALB : from Albania and G1, G2, G3: from Greece)
- ✓ *Bornmuellera tymphaea* (BORN)
- ✓ *Leptoplax emarginata* (LEPT)
- ✓ *Nocca tymphaea* (NOC)



Cropping

70 days



Harvest




Analyses

- Dry biomass weight
- Metal concentrations (ICP)
- (Al, As, B, Cd, Co, Cr, Cu, Fe, Mn, Ni, Pb, Zn and Ca, K, Mg, Na, P)


Results

*Error bars represent standard errors

Biomass of shoots of the seven populations *



Ni Concentration in shoots of each population *



❖ *L. emarginata* has the highest biomass (mean: 0.38 g) whereas the highest Ni concentration has been recorded in *B. tymphaea* (mean: 3 956 mg kg⁻¹) (Wilcoxon, p<0.05).

❖ The 4 populations of *A. murale* have led to different results, the best one being *A. murale* ALB.

❖ In terms of Ni quantity, *L. emarginata*, *B. tymphaea* and *A. murale* ALB (results not show here) are the more efficient populations.



Conclusions

The intra- and inter- population variabilities of Ni accumulation in the four species are very high. This screening enables us to retain *L. emarginata*, *B. tymphaea* and *A. murale* ALB as the most promising populations for Ni phytoextraction.

Furthermore, the individuals accumulating the highest mass of Ni also concentrate Ca, Mn, and Co.

However, these results have been recorded for 70 d aged individuals. Elemental concentrations depend on the phenological stage. Indeed, the measured metal concentrations are lower expected for individuals collected in the field at the flowering stage.

A principal component analysis has been run with all the metal concentrations above 10 mg_{metal} (kg_{dry matter})⁻¹. Therefore, Cr, Cu, Cd and Na have been neglected (however it was noticed that Cd concentration was 18 times higher in *N. tymphaea* than in the other ones).



AG1, AG2, AG3 and ALB: four populations of *A. murale*, NOC: *N. tymphaea*; BORN: *B. tymphaea*; LEPT: *L. emarginata*.

❖ Concentrations of Ni, P, Ca, Co Mn and B are correlated. At the same time, Zn, Fe, Al, Mg and K concentrations are correlated but not with the previous group.

❖ Biomass production is opposed to concentration of Zn, Mg, Fe, K and Al in shoots.

Perspectives

This study brings criteria to select the optimal population and individuals for further hydrometallurgical process.

This work could be extended to multicontaminated matrices, including tolerance to a co-contamination as a screening factor of hyperaccumulators.