



Intercropping legume and non-legume, an innovative way to valorize N₂ fixation and soil mineral sources in low inputs cropping systems.

Laurent Bedoussac, Etienne-Pascal Journet, Henrik Hauggaard-Nielsen, Christophe Naudin, Guenaelle Corre - Hellou, Loïc Prieur, Erik Steen Jensen, Eric E. Justes

► To cite this version:

Laurent Bedoussac, Etienne-Pascal Journet, Henrik Hauggaard-Nielsen, Christophe Naudin, Guenaelle Corre - Hellou, et al.. Intercropping legume and non-legume, an innovative way to valorize N₂ fixation and soil mineral sources in low inputs cropping systems.. 18. Nitrogen workshop "The nitrogen challenge: building a blueprint for nitrogen use efficiency and food security", Jun 2014, Lisbonne, Portugal. pp.573, 10.13140/2.1.3836.2249 . hal-02741691

HAL Id: hal-02741691

<https://hal.inrae.fr/hal-02741691>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



INTERCROPPING LEGUME AND NON- LEGUME, AN INNOVATIVE WAY TO VALORIZE N₂ FIXATION AND SOIL MINERAL N SOURCES IN LOW INPUTS CROPPING SYSTEMS


Bedoussac L., Journet E.-P., Hauggaard-Nielsen H.,
Naudin C., Corre-Hellou G., Prieur L., Jensen E.S., Justes E.



Intercrops (or mixed crops)

Simultaneous growth of two or more species in the same field for a significant period without necessarily sowing and harvesting them together (Willey 1979)

- Application of ecology principles like biodiversity and species interactions (e.g. Vendermeer, 1989)
- Better use of natural resources in time and space



Separated rows



OR

Mixed on the row





Material and objectives

- ❑ 48 organic experiments from 2001 to 2010
- ❑ 3 pedoclimatic situations
- ❑ Spring and Winter crops





- ❑ **Large range of practices** : cultivars, densities, organic fertilization
- ❑ **Aim** :
 - ❑ Evaluate the potential advantages of intercrops for grain yield, grain protein concentration and weed control
 - ❑ Analyze the functioning of cereal-grain legume intercrops to further propose optimized intercropped systems.



Species complementarity

for N sources (soil mineral N and N₂ from air)




- Wheat roots deeper than those of the legume
→ **Deep nutrients only available for the cereal**

- Higher legume N₂ fixation rate in IC (75% vs. 62%)
→ **Most of soil N mineral available for the cereal**



IC improve yield (compared to low N sole crops)

(Hauggaard-Nielsen et al. 2001; Bedoussac and Justes 2010)




- Total intercrop grain yield higher than the mean sole crops (3.3 vs 2.7 Mg ha⁻¹)
→ **Highest efficiency for low N**
- Total IC grain yield more stable compared to each sole crop
→ **Higher resiliency**
- Proportion of cereal > 50%
→ **Cereal more competitive**



IC improve grain quality

(Jensen 1996; Hauggaard-Nielsen et al. 2001, 2009; Bedoussac and Justes 2010)




- Cereal grain protein concentration higher in IC (11.1% vs. 9.8% in SC)
- The lowest the SC protein the higher the increase
→ **Highest efficiency for low N**
- Due to :
 - lower cereal grain yield
 - low use of soil mineral N by the legume (75% of Ndfa)
→ **More N available per plant, tiller & grain in IC**



IC reduce weeds (in comparison of legume)

(Hauggaard-Nielsen et al. 2001, Corre-Hellou et al. 2011)



- Less weeds in IC compared to the legume (0.40 vs. 1.38 Mg ha⁻¹)
- No difference compared to the SC cereal

→ **In the IC weeds mostly controlled by the cereal**

→ **Less light and N available for the weeds**



Conclusion and perspectives

- ❑ **Intercropping is an efficient way to improve yield, quality and reduce weeds in low inputs systems**
- ❑ **Intercropping development need the collaboration of all the actors** in the value chain (farmers, collectors, breeders, agribusiness companies...)
- ❑ **Modelling intercropping systems could be helpful** to optimize them and to determine varietal characteristics suited to mixtures





Obrigado pela sua atenção

Bedoussac L., Justes E., Journet E.-P., Hauggaard-Nielsen H., Naudin C., Corre-Hellou G., Prieur L., Jensen E. S.

Intercropping, an application of ecological principles to improve nitrogen use efficiency in organic farming systems

In: Organic farming, prototype for sustainable agricultures

Bellon S. et Penvern S. (eds), Springer, Berlin (2013)



Technical University
of Denmark



Nourrir le monde d'intelligences