



HAL
open science

Regeneration of the rat tibialis anterior muscle is impaired despite induction of the SPARC- β -catenin pathway during post-immobilization recovery

Lamia Slimani, Emilie Vazeille, Christiane Deval, Julien Amat, Cécile Polge, Daniel D. Bechet, Daniel Taillandier, Dominique Dardevet, Anne Listrat, Didier Attaix, et al.

► To cite this version:

Lamia Slimani, Emilie Vazeille, Christiane Deval, Julien Amat, Cécile Polge, et al.. Regeneration of the rat tibialis anterior muscle is impaired despite induction of the SPARC- β -catenin pathway during post-immobilization recovery. *Molecular Biology of Muscle Development and Regeneration*, May 2014, Lecce, Italy. , 2014, *Molecular Biology of Muscle Development and Regeneration*. hal-02741741

HAL Id: hal-02741741

<https://hal.inrae.fr/hal-02741741v1>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Background & Aims

The immobilization-induced tibialis anterior (TA) muscle atrophy worsens after cast removal concomitantly with changes in the extracellular matrix composition^{1,2}.


SPARC is a matricellular glycoprotein involved in tissue response to injury and in stabilization of β -catenin, which induces muscle regulatory factors (MRFs) controlling muscle regeneration.

We hypothesized that SPARC expression changed upon immobilization and could be involved in the worsening of TA muscle atrophy by altering muscle regeneration processes pending cast removal.

¹Vazeille et al. Am J Physiol Endocrinol Metab 2008, ²Slimani et al. Am J Physiol Endocrinol Metab 2012

Methods


Wistar rats were subjected to unilateral hindlimb immobilization for 8 days (I8) or not (I0), and allowed to recover for 1 to 10 days (R1-10).




Expression of SPARC, β -catenin, and proliferative (i.e. MyoD and Myf5) or differentiation (i.e. myogenin) MRFs were assessed by Western blots and/or RT-qPCR in previously immobilized TA during recovery.

I/ Tibialis anterior muscle atrophy worsened rapidly after cast removal concomitantly with endomysium thickening

Muscle atrophy




Tibialis anterior muscle atrophy is expressed as % difference vs. controlateral control leg are means \pm SEM (n=10-11 rats/group). Bars with different letters are significantly different.




Ten-micrometer-thick cross-sections from control (CTL) and immobilized (IMM) tibialis anterior muscles were stained with Picro-Sirius red, which reveals intramuscular connective tissue. Observations and image acquisitions were performed using a photonic microscope in bright-field mode (Olympus BX-51, Tokyo, Japan) coupled to a high-resolution cooled digital camera (Olympus DP72) and Cell-D software (Olympus Soft Imaging Solutions, Münster, Germany). E, endomysium; P, perimysium.

After quantification, endomysium and perimysium areas were expressed as % of total field and are means \pm SEM (n=5-6 rats/group). Dots with different letters are significantly different.





II/ Sparc mRNA levels also increased rapidly after cast removal






mRNA levels in control and immobilized tibialis anterior muscles were assessed by RT-qPCR, expressed as fold induction vs. I0 group and are means \pm SEM (n=10-11 rats/group). Bars with different letters are significantly different.

III/ β -catenin mRNA and protein levels increased during immobilization and/or recovery in a time dependent manner



β -catenin protein levels from control (Ctl) or immobilized (Imm) tibialis anterior muscles were detected using a polyclonal antibody (Sigma-Aldrich). Signals were quantified after acquisition using a chemiluminescence imaging system (Syngene G:BOX XT4). Data are arbitrary units, were normalized against ponceau red staining for uneven loading, and are means \pm SEM (n=10-11 rats/group). Bars with different letters are significantly different.

IV/ MyoD and Myf5 mRNA levels increased concomitantly to increased β -catenin expression at I8 and R1, whilst myogenin mRNA levels first decreased at I8 and R1, and increased thereafter concomitantly to increased β -catenin expression at R10.



Conclusions

We report an induction of the SPARC- β -catenin pathway associated with increased mRNAs of the proliferative MRFs (Myf5 and MyoD) in the recovering TA early after cast removal. The differentiation MRF myogenin was first largely repressed, but increased later on, when TA started to recover. Altogether, the data suggest that the TA tended to preserve muscle regeneration potential through induction of proliferative MRFs. However this process was poorly efficient presumably because of an alteration myogenic differentiation.