

Valorisation of the natural variation available in tomato accessions to decipher the genetic control of fruit quality traits

Laura Pascual-Banuls, Jiaxin Xu, Christopher Sauvage, Guillaume Bauchet, Nelly Desplat, Jean-Paul Bouchet, Dominique D. Brunel, Mathilde M. Causse

▶ To cite this version:

Laura Pascual-Banuls, Jiaxin Xu, Christopher Sauvage, Guillaume Bauchet, Nelly Desplat, et al.. Valorisation of the natural variation available in tomato accessions to decipher the genetic control of fruit quality traits. Applied Vegetable Genomics, Feb 2014, Vienne, Austria. hal-02741808

HAL Id: hal-02741808 https://hal.inrae.fr/hal-02741808

Submitted on 3 Jun2020


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





International Conference

Applied Vegetable Genomics



Valorisation of the Natural Variation Available in Tomato Accessions to Decipher the Genetic Control of Fruit Quality Traits

L. Pascual¹, J. Xu^{1, 2}, C. Sauvage¹, G. Bauchet¹, N. Desplat¹, J. Bouchet¹, D. Brunel³, M. Lepaslier³, <u>M. Causse¹</u>

¹ INRA, UR1052, GAFL, BP94,F-84143 Montfavet, France (mathilde.causse@avignon.inra.fr), ² NorthwestA&FUniversity, College of Horticulture, Yang Ling, Shaanxin, 712100, P.R.China, ³ INRA, EPGV, Centre de Génotypage, Evry, France

Identifying the genes controlling the variation of quantitative traits is a key goal for breeders. Genetic variations underlying quantitative traits (QTL) have been mapped by traditional linkage mapping for years and positional cloning identified several QTLs. However linkage mapping is limited to the analysis of traits differing between two lines and the impact of genetic background on QTL effect has been underlined. Thanks to the increase in molecular markers, genome-wide association studies were then proposed to circumvent QTL limitations. In tomato, a self-pollinated crop, we have shown that association studies are possible, using the admixed nature of cherry tomato genomes that limits the impact of population structure in such an approach. Nevertheless, the results might be limited by linkage disequilibrium, which varies greatly across the genome. Multi-allelic Advanced Generation Inter-Cross (MAGIC) populations allow a wide range of variability to be analyzed and avoid dealing with population structure.

We have constructed a MAGIC population by crossing 8 tomato lines, representing a wide range of genetic diversity within the species. The large range of phenotypic variability represented by these lines was then characterized at different scales, metabolomic, proteomic and transcriptomic. The variation of a few genes/proteins, involved in primary metabolism and stress response, was related to metabolite content by network reconstruction. The whole genomes of the 8 founder lines were resequenced identifying more than 4 millions SNPs when mapped onto the reference genome. A set of 1536 SNPs markers was then selected to genotype the MAGIC population and used to construct a linkage map. A large increase in recombination frequencies compared to bi-parental populations was shown. QTLs for fruit quality traits were mapped and related to the variations detected at the genome sequence and expression levels in the parental lines. The advantages and limits of the three types of population, in the context of the available genome sequence and resequencing facilities, will be discussed.