



HAL
open science

Amino acids as orobanchicides: an innovative approach for biocontrol broomrape weeds.

Monica Fernandez-Aparicio, Stéphanie Gibot-Leclerc, Delphine Moreau, Christian Steinberg, David Bohan, Bruno Chauvel, Alexandre Bernard, Pascal Marget, Laurent Falchetto, Pascal Farcy, et al.

► **To cite this version:**

Monica Fernandez-Aparicio, Stéphanie Gibot-Leclerc, Delphine Moreau, Christian Steinberg, David Bohan, et al.. Amino acids as orobanchicides: an innovative approach for biocontrol broomrape weeds.. 17. European weed research society symposium, Jun 2015, Montpellier, France. , 2015, Proceedings of the 17th European Weed Research Society Symposium. hal-02741844

HAL Id: hal-02741844

<https://hal.inrae.fr/hal-02741844v1>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



17th EUROPEAN WEED RESEARCH SOCIETY SYMPOSIUM **EWRS 2015**

23-26 June 2015



PROCEEDINGS

Montpellier SupAgro, France



AMINO ACIDS AS OROBANCHICIDES: AN INNOVATIVE APPROACH FOR BIOCONTROL OF BROOMRAPE WEEDS

Fernandez-Aparicio M.¹, Gibot-Leclerc S.², Moreau D.¹, Steinberg C.¹, Bohan D.A.¹, Chauvel B.¹, Bernard A.³, Marget P.³, Falchetto L.³, Farcy P.³, Ratchinskiy C.¹, Gerbaud B.¹, Boari A.⁴, Vurro M.⁴, Sands D.C.⁵, Morris C.E.⁶, Buttin P.⁷, Arhancet G.B.⁷, Reboud X.¹

¹ INRA, UMR1347 Agroécologie, Dijon, France ; ² Agrosup-Dijon, UMR1347 Agroécologie, Dijon, France ; ³ INRA, UE 0115 Domaine Experimentale d'Epoisses,-Brenetiere, France ; ⁴ CNR, Bari, Italy ; ⁵ Montana State University, Bozeman, USA ; ⁶ INRA, UR0407 Pathologie Vegetale, Montfavet, France ; ⁷ Novus International Inc, St Charles, USA
monica.fernandez@dijon.inra.fr

Broomrapes (*Orobanche* and *Phelipanche* spp) are obligate, chlorophyll-lacking, root parasitic weeds that obtain all their nutrient resources by feeding off many important crops via their haustoria. France suffers from severe infestations of three broomrape species, *O. cumana* Wallr. infecting sunflower, *O. minor* Sm. infecting forage legumes and *P. ramosa* L. (syn *O. ramosa*) infecting hemp, oilseed rape, and tobacco. Several broomrape traits such as mass seed production, their easy dispersal and longevity, germination induced by host root exudates, specialized host range, and physical and metabolic overlap with the host make the management of these weeds very difficult. Successful control strategies must target parasitic subterranean stages in order to hamper both the main parasitic sink for nutrients and the subsequent parasitic seed production. An effective strategy to control broomrape is the foliar application of systemic herbicides to herbicide-resistant crops. The crop will deliver the herbicidal effect into the broomrape attached to their roots by translocating the herbicide systemically to the young parasite via the haustorium at the host-parasite interface. Among those herbicides, glyphosate, imidazolinones and sulfonylureas specifically inhibit either the aromatic or the branched-chain amino acid synthesis in the parasite by targeting key enzymes in their synthesis pathways. This method is one of the few available strategies efficient for broomrape control however it presents two main drawbacks. First, herbicide resistance is not available for many important crops and second, increasing concerns for environmental pollution. An environmental-friendly alternative to this method could be the use of broomrape-specific inhibitory amino acids. Amino acid synthesis pathways are regulated by feedback inhibition of controlling enzymes by some of the amino acid end-products resulting in starvation of other amino acids in the same pathway and as consequence growth inhibition. The suppressive activity of amino acid end-products varies according with species and life stages. Laboratory experiments revealed that methionine and lysine inhibit the germination and radicle elongation of the broomrape species affecting French agriculture. This developmental inhibition hampers the infective potential of broomrape seedbank in the host crop. Additional mechanisms such as elicitation of resistance by methionine against *Orobanche* root penetration as well as the use of selected and characterized strains of *Fusarium oxysporum* f. sp. *orthoceras* are being investigated. Methionine and lysine are produced in large scale as animal feed supplements and can be delivered to the soil during the short infective period in which broomrape germination and host infection occur. The persistence of the amino acid is predicted to be low beyond this short time frame due to soil microbial metabolism. The *Orobanche* Biocontrol Project at INRA-Dijon is carrying out in vitro, rhizotron and field experiments aimed to optimize the potential use of methionine and lysine as orobanchicides. This project is funded by Santé des Plantes et Environnement division (SPE-www.spe.inra.fr).