Does biostimulation of grapevine impact elicitor-induced resistance against downy mildew ? A methodological framework

Yuko Krzyzaniak, Marie-Claire Heloir, Sophie Trouvelot, Estelle Moreau, Marielle Adrian

Does biostimulation of grapevine impact elicitor-induced resistance against downy mildew? A methodological framework

Krzyzaniak Yuko [1], Héloir Marie-Claire [1], Moreau Estelle [2], Trouvelot Sophie [1], Adrian Marielle [1], and other members of IRIS+ consortium

(1) Agroécologie, AgroSup Dijon, CNRS, INRA, Univ. Bourgogne Franche-Comté, F-21000 Dijon, France
(2) Parc Technopolitain Atalante, CS41908 35435 Saint Malo cedex, France

yuko.krzyzaniak@dijon.inra.fr
+33 (0)3 80 69 34 15
17 rue Sully - 21065 Dijon FRANCE

Context

Biostimulants (BS) are applied to crops in order to improve [1] :

- the yield and quality of harvested organs
- the uptake of nutrients
- tolerance to abiotic stresses

BUT …*How do they act and how to assess it ?*

We aim at deepening the knowledge of their **mode of action on grapevine** (*Vitis vinifera, L.), a p*erennial crop of economic and cultural value.

MOREOVER…

Grapevine is susceptible to cryptogamic diseases such as **downy mildew** caused by *Plasmopara viticola.*

Protection strategies using **resistance inducers** (RIs) on those crops are well documented and validated in controlled conditions[2]. However, the efficacy of RIs is **variable** in field conditions, and may **depend on the physiological status** of the plant [3, 4, 5].

HYPOTHESIS

An improvement of the plants' physiological status by biostimulation is expected to increase their responsiveness to RI application.



Aims and global approach of the IRIS+ project

1 **To define and develop a panel of tools and methods** to study the impact of biostimulants on the development and the physiology of grapevine in greenhouse conditions.

2 **To test whether biostimulation can increase** grapevine responsiveness to RI application, *via* an improvement of its physiological status.

BS application(s) → Analysis of the effects on plants → RI application → Inoculation with the pathogen → Increased protection rate compared to RI only ?

Foliar and/or root applications of BS (or water for control) on potted grapevine herbaceous cuttings (cv Marselan) in greenhouse conditions.

Measurement the impacts of BS on grapevine development and physiology, compared to water-treated control.

Foliar application of RI (or water), 2 days before inoculation with Plasmopara viticola.

Assessment of the efficacy of RI by measuring the sporulating leaf area.

Development of tools and methodology

A Definition of culture systems and growth conditions

Pot system

Rhizotron system

B Evaluation of the effects of biostimulation

PLANT – ORGAN scale

Aerial and root phenotyping
Ex: Non destructive growth and development tracking devices.

Photosynthetic activity
Ex: Photosynthesis analysis by gas exchange measurements (CO_2, H_2O).

TISSUE – CELLULAR scale

Chlorophyll fluorescence
Ex: Imaging-PAM. Image capture and analysis for quantifying chlorophyll fluorescence of leaves.

Defence events
Ex: Vizualisation of H_2O_2 production sites (diaminobenzidine staining).

MOLECULAR scale



Gene expression and related enzymatic activity
Ex: Targeted analysis of the expression level of genes involved in physiology/defence (encoding for invertases, PR-proteins, amongst others).



Metabolic profiles
Ex: Analysis of primary and secondary metabolites in response to treatments (carbohydrates, phytoalexins, phytohormones).

C Evaluation of elicitor-induced resistance



Protection rate
Ex: Image analysis of sporulating area of leaf disks infected by *P. viticola*.

[1] European Biostimulants Industry Consortium, EBIC (2012) The 1st World Congress on the use of Biostimulants in Agriculture- Strasbourg Congress Center, France
[2] Delaunois *et al*. (2014). Elicitors as alternative strategy to pesticides in grapevine? Current knowledge on their mode of action from controlled conditions to vineyard. *Environ Sci Pollut Res* (2014) 21:4837–4846
[3] Bolton MD.(2009.) Primary metabolism and plant defense-fuel for the fire. *Mol Plant Microbe Interac,* 22: 487–497
[4] Dietrich *et al*. (2005). Growth responses and fitness costs after induction of pathogen resistance depend on environmental conditions. *Plant, Cell and Environment* 28: 211–222
[5] Maymoune A. *et al*. (2015). Impact of abiotic stresses on the protection efficacy of defence elicitors and on metabolic regulation in tomato leaves infected by *Botrytis cinerea*. *Eur J Plant Pathol*, 142: 223-237