



HAL
open science

Identification of two new genes involved in myocyte fusion in vertebrates

Aurélie Landemaine, Jérôme Bugeon, Pierre-Yves Rescan, Jean-Charles Gabillard

► **To cite this version:**

Aurélie Landemaine, Jérôme Bugeon, Pierre-Yves Rescan, Jean-Charles Gabillard. Identification of two new genes involved in myocyte fusion in vertebrates. 9. European Zebrafish Meeting, Jun 2015, Oslo, Norway. , 2015, Zebrafish 2015. hal-02742363

HAL Id: hal-02742363

<https://hal.inrae.fr/hal-02742363v1>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



FINAL PROGRAMME

The 9th European Zebrafish Meeting

Oslo, Norway, June 28-July 2, 2015



[P1-061] Identification Of Two New Genes Involved In Myocyte Fusion In Vertebrates

Aurélie Landemaine [FRANCE]¹, Jérôme Bugeon [FRANCE]¹, Pierre-Yves Rescan [FRANCE]¹, Jean-Charles Gabillard [FRANCE]¹

INRA-LPGP¹

In most animals, myocyte fusion is a cellular process highly regulated during skeletal muscle development, post embryonic muscle growth and regeneration. Myocyte fusion follows an ordered set of cellular events, including myocyte elongation, cell migration, recognition and adhesion, and membrane fusion. However, our knowledge regarding molecular mechanisms controlling this essential step in myogenesis is far from being complete. Our study aimed at identifying new genes involved in myocyte fusion step in vertebrates. First, we performed an in vitro functional screening in C2C12 cell line (murine model) based on siRNA knock-down. Using a selection of genes found to be expressed in proliferating and differentiating muscle C2C12 cells (Moran and al., 2002 ; Tomczak and al., 2004), our functional screening led to the identification of thirteen genes which in vitro knock-down induced a myocyte fusion default. To identify their roles in vivo, expressional and functional analysis of these genes was further conducted on zebrafish model. The expression of four genes out of thirteen was clearly seen in somite during embryonic myogenesis using whole mount in situ hybridization. To precise their functions, in vivo studies based on morpholino micro-injection were performed on zebrafish embryos. Knock-down of two candidate genes, one encoding a protein of the extracellular matrix and the other encoding a N-acetyltransferase enzyme, resulted in an inability of the injected embryo to swim after hatching. We hypothesize that a muscle structural disorganization was responsible of this functional default, a confocal observation is currently in progress to characterize the muscle structure.
