



HAL
open science

Dynamic and structural studies of lipid droplets using synchrotron light

Frederic Jamme, Jean-David J.-D. Vindigni, Alexandre A. Giuliani, Franck Wien, Roselyne Tâche, Franjo Jagic, Mathieu Réfrégiers, Pascale P. Jolivet, Thierry Chardot, Yann Gohon, et al.

► To cite this version:

Frederic Jamme, Jean-David J.-D. Vindigni, Alexandre A. Giuliani, Franck Wien, Roselyne Tâche, et al.. Dynamic and structural studies of lipid droplets using synchrotron light. Conférence Jacques Monod, Analyses Moléculaires de l'Organisation et du Remodelage des Membranes, Nov 2014, Roscoff, France. 2014. hal-02742432

HAL Id: hal-02742432

<https://hal.inrae.fr/hal-02742432>

Submitted on 3 Jun 2020


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CONTEXT

Lipid droplet: a complex and dynamic organelle

In cells, neutral lipids (triglycerides and steryl esters) are stored in organelles called lipid droplets (LDs) [1]. They are present in all organisms, from bacteria to plants and animals.



Lipid droplets: not well known but with rising interest

From biologists

LD is not an inert fat depot but a dynamic organelle which regulates cell metabolism and signaling

From medical field

LDs have a crucial role in diseases with increasing prevalence (obesity, diabetes) [2]
Oleosins (from peanut and hazelnut), seed LD associated proteins are allergens [3].

From industrials

crushing: oils for food and non food productions are extracted from seed LDs
food processing industry, cosmetic and health: oleosins harbor interfacial properties and could be used as emulsifying agents or in drug delivery systems [4]




Oleosins, seed lipid droplet associated proteins

Oleosins, AtOle1 and AtClo1, are LD integral proteins

Predicted structure = tri-block organization [5]:

- variable N-terminal and C-terminal part, exposed at the surface and in contact with the cytosol
- highly hydrophobic central part inserted into the phospholipid (PL) monolayer and/or the triacylglycerol (TG) core.




Questions and objectives

- Role on lipid filling
- Role on LD structure and stabilization
- Structural data on oleosins inserted into LD (natural environment)

HETEROLOGOUS EXPRESSION OF PLANT OLEOSINS IN YEAST


Oleosins are targeted to LDs in yeast

Photonic microscopy pictures (bright field and epifluorescence) of yeast expressing AtClo1-GFP [6].




Ole1 and Clo1 become the major proteins associated with yeast LDs

SDS-PAGE protein profiles of LDs purified on sucrose gradient




Oleosins induce LDs and lipid accumulation

Thin sections of yeasts expressing AtOle1-GFP (transmitted electron microscopy)



LDs, round and white structures, are accumulated in oleosin expressing cells





Lipid classes were analyzed using thin layer chromatography after Folch extraction




oleosins induce neutral lipid accumulation

DYNAMIC STUDY USING SYNCHROTRON FTIR

Single cell FTIR analysis on Soleil SMIS beamline revealed a link between neutral lipid and carbohydrate fluxes [7].



Metabolic modifications were confirmed using biochemical analysis





More neutral lipids triacylglycerol and steryl esters
Less storage carbohydrates trehalose and glycogen

STRUCTURAL STUDY USING SRCD



SRCD at DISCO beamline revealed that Ole1 is mainly beta folded when inserted in LDs [9]

Summary of oleosin structural studies



LDs purified on sucrose gradient and observed using microscopy

First SRCD spectra on purified organelle



Hydrophobic domain of oleosins = beta sheet hairpin


Perspectives on LD study

Lipid core organization?

- Small-angle X-ray scattering SAXS (SWING)
- Membrane fluidity using fluorescence anisotropy (DISCO)
- Protein mapping using Deep UV (DISCO)
- 3D imaging with UV (DISCO)

LD and other cell compartments?

- 3D imaging with UV (DISCO)



Other oleosins

- SRCD on Ole3 in LDs (DISCO)
- Synchrotron water radiolysis footprinting of accessible amino-acids (METROLOGY)

Protein with hairpin hydrophobic core

- Stomatol in progress, caveolin?, GPAT4? HCV core protein?

Other LD proteins

- Perilipins, Apolipoproteins

