



HAL
open science

Analysis of time-series of total and plant water stress levels using a dual-source energy balance model over agricultural crops and medium to low resolution thermal infra red remote sensing data

Gilles Boulet, Bernard Mougenot, Malik Bahir, Pascal Fanise, Sameh Saadi, Vincent Simonneaux, Wafa Chebbi, Zeineb Kassouk, Toufik Oualid, Albert Oliosio, et al.

► **To cite this version:**

Gilles Boulet, Bernard Mougenot, Malik Bahir, Pascal Fanise, Sameh Saadi, et al.. Analysis of time-series of total and plant water stress levels using a dual-source energy balance model over agricultural crops and medium to low resolution thermal infra red remote sensing data. EGU 2015, European Geosciences Union General Assembly, European Geosciences Union (EGU). DEU., Apr 2015, Vienne, Austria. hal-02742459

HAL Id: hal-02742459

<https://hal.inrae.fr/hal-02742459>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Analysis of time-series of total and plant water stress levels using a dual-source energy balance model over agricultural crops and medium to low resolution thermal infra red remote sensing data

Gilles Boulet (1,2), Bernard Mougnot (1,2), Malik Bahir (3), Pascal Fanise (1), Sameh Saadi (1,2), Vincent Simonneaux (1), Wafa Chebbi (2), Zeineb Kassouk (2), Toufik Oualid (4), Albert Olioso (5), Jean-Pierre Lagouarde (6), Valérie Le Dantec (7), Vincent Rivalland (8), Mehrez Zribi (8), Zohra Lili-Chabaane (2,9)

(1) IRD, UMR5126 CESBIO, Toulouse, France (gilles.boulet@ird.fr), (2) Institut National Agronomique de Tunisie, Tunis, Tunisia, (3) CNES/CNRS, UMR5126 CESBIO, Toulouse, France, (4) LRSBG, Université de Mascara, Mascara, Algeria, (5) INRA, UMR1114 EMMAH, Avignon, France, (6) INRA, UMR1391 ISPA, Villenave d'Ornon, France, (7) Université de Toulouse, UMR5126 CESBIO, Toulouse, France, (8) CNRS, UMR5126 CESBIO, Toulouse, France, (9) Université de Carthage, Tunis, Tunisia

Detecting, monitoring and mapping plant water stress with remote sensing data is a crucial component of modern agricultural water management, especially in areas with scarce water resources such as the south and the eastern parts of the Mediterranean region. Developing efficient operational methods dedicated to those three actions is thus necessary to design observing systems for areas with a mixture of irrigated, rainfed and deficit irrigation agriculture. Those systems can assist managers in tasks such as early warning of drought, real time irrigated area mapping etc. A way to quantify plant and total water stress levels is to exploit the available surface temperature data from remote sensing as a signature of the surface energy balance, including the latent heat flux. Remotely sensed energy balance models enable to estimate evapotranspiration and the water status of continental surfaces. Two-source models, such as TSEB (Norman et al., 1995) allow deriving a rough estimate of the water stress of the vegetation instead of that of a soil-vegetation composite. For the latter, a realistic underlying assumption enables to invert two unknowns (evaporation and transpiration) from a single piece of information. This assumption states that, in most cases, vegetation is unstressed, and that if vegetation is stressed, evaporation is negligible. In the latter case, if vegetation stress is not properly accounted for, the resulting evaporation will decrease to unrealistic levels (negative fluxes) in order to maintain the same total surface temperature. Actual and potential transpiration rates are combined to derive an index of plant water stress applicable to low resolution data. Here, we evaluate time series of plant water stress indices in the Kairouan area in Central Tunisia in the last few years by comparing them with 1- maps of the irrigation sectors as well as rainfall data and 2- turbulent heat flux measurements obtained at low resolution (scintillometer, eddy-covariance over homogeneous areas) and 3- outputs of a distributed hydrological model (SAMIR).