



**HAL**  
open science

## Can arbuscular mycorrhizal fungi be used as bioindicators of land use ?

Marie-Lara Bouffaud, C. Bragalini, Diederik van Tuinen, S. Voyron, E.  
Lumini, Daniel D. Wipf, V. Bianciotto, M. Girlanda, Dirk Redecker

### ► To cite this version:

Marie-Lara Bouffaud, C. Bragalini, Diederik van Tuinen, S. Voyron, E. Lumini, et al.. Can arbuscular mycorrhizal fungi be used as bioindicators of land use?. 1. Global Soil Biodiversity Initiative Conference (GSBI), Institut National de la Recherche Agronomique (INRA). Dijon, FRA., Dec 2014, Dijon, France. 705 p. hal-02742502

**HAL Id: hal-02742502**

**<https://hal.inrae.fr/hal-02742502>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THE FIRST GLOBAL  
**SOIL BIODIVERSITY CONFERENCE**

Assessing soil biodiversity and role in ecosystem services



GLOBAL  
SOIL BIODIVERSITY  
INITIATIVE

Palais des Congrès, Dijon, France

**2-5 DECEMBER 2014**

THE FIRST GLOBAL  
**SOIL BIODIVERSITY**  
**CONFERENCE**

December 2-5, 2014, Dijon, France

Book of Abstracts

# PROGRAM AND ABSTRACTS

## THE FIRST GLOBAL **SOIL BIODIVERSITY** **CONFERENCE**



GLOBAL  
**SOIL BIODIVERSITY**  
INITIATIVE

December 2-5, 2014

Dijon – France

[P2.238]

**Can arbuscular mycorrhizal fungi be used as bioindicators of land use?**

M.L. Bouffaud<sup>1</sup>, C. Bragalini<sup>2,3</sup>, D. Van Tuinen<sup>1</sup>, S. Voyron<sup>2</sup>, E. Lumini<sup>4</sup>, D. Wipf<sup>1</sup>, V. Bianciotto<sup>4</sup>, M. Girlanda<sup>2</sup>, D. Redecker<sup>1</sup>

<sup>1</sup>INRA Dijon, France, <sup>2</sup>University of Turin, Italy, <sup>3</sup>Université Claude Bernard Lyon1, France, <sup>4</sup>National Research Council (CNR) Turin, Italy

Being abundant in nearly all soils and less diverse than other soil organisms, Arbuscular Mycorrhizal Fungi (AMF) are potential indicators of land management legacies and soil quality degradation. It has been pointed out that these are critical factors for understanding and supporting the sustainable use of soils, but can be difficult to measure directly (Jansa et al. 2014). To serve as broadly applicable bioindicators, AMF should exhibit consistent patterns in a range of soils.

In the framework of the EcoFINDERS project, soil AMF assemblages were described in four European long term observatories (mainly grasslands), located in different climatic and geological zones, subjected to either low or high management intensity. AMF communities were described, in both spring and autumn, by means of high-throughput metabarcoding targeting the ITS2 region. AMF community structures were related to soil properties, land management intensity and geographic distances to address the relative importance of these factors in shaping the composition of AMF communities.

As expected, our results indicate significant effects of management type ( $p=0.000999$ ), soil chemical and physical properties (such as pH, soil texture, organic carbon content and available phosphorus), and among-site geographic distances on community composition (Mantel tests). Indicator species analysis (presence/absence data) retrieved some taxa (or their combinations) characteristic of specific land uses (or their combinations) in both seasons.

Jansa J. et al. 2014 - Soil and geography are more important determinants of indigenous arbuscular mycorrhizal communities than management practices in Swiss agricultural soils. *Molecular Ecology* 23: 2118–2135

Keywords: Glomeromycota, bioindicators, land use