



Nitrogen use efficiency in bread wheat: breeding tools and recent genetic progress

Fabien Cormier, Sébastien Faure, Pierre Dubreuil, Emmanuel E. Heumez,
Katia Beauchêne

► To cite this version:

Fabien Cormier, Sébastien Faure, Pierre Dubreuil, Emmanuel E. Heumez, Katia Beauchêne. Nitrogen use efficiency in bread wheat: breeding tools and recent genetic progress. Wheat Breeding 2014: Tools, targets and progress, Association of Applied Biologists. Warwick, GBR., Jan 2014, Harpenden, United Kingdom. hal-02742524


HAL Id: hal-02742524

<https://hal.inrae.fr/hal-02742524v1>

Submitted on 3 Jun 2020


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.




Nitrogen Use Efficiency in Bread Wheat: breeding tools and recent genetic progress

Fabien Cormier, Sébastien Faure, Pierre Dubreuil, Emmanuel Heumez, Katia Beauchêne,
Stéphane Lafarge, Sébastien Praud, Jacques Le Gouis




What are the N traits to select for ?

- **GPD** = Grain Protein Deviation
 - Proposed by Monaghan et al (2001)
 - Bonus for GPD+ lines during registration in France (2007)



- **NUE** = Nitrogen Use Efficiency ?
 - First official trials at 3 N levels harvested in 2013
 - Identification of potential criteria after two experimental years
 - Bonus for « efficient » lines by 2016 ?



What is Nitrogen Use Efficiency?

- Two main components

✓ **N Uptake** = total plant N / available N

✓ **N Utilization** = grain yield / total plant N


and sub-components : Straw N content, Grain N content, ...

- Two main possible targets




✓ **Sub-optimal** -> Maintain yield / quality

✓ **Optimal** -> Increase yield / quality

that will depend on Genotype x N level interactions



Can we improve NUE?



NUE = GY if we suppose Available N to be the same for all genotypes

What about recent breeding progress on NUE ?

Has NUE been increased ? How? For which N levels?

What are traits heritabilities?

Is there GxN interactions?

In which environment to select?



Is elite diversity sufficient?

A panel of registered lines

225 European elite varieties

160 "French" + 65 others

Registered from 1985 to 2010



A network of field experiments

Site x Season	Years	Genotypes tested	Residual soil N (kg N ha ⁻¹)	N supply ¹ (kg ha ⁻¹)	
				HN	LN
VB08	07/08	197	106	126	44
VR09	08/09	196	30	220	120
EM08	07/08	207	67	170	70
EM09	08/09	208	30	150	50


8 trials = 2 N x 4 environments

with environmental characterization (daily temperature, rain,...)

2865 plots - 10 measured traits

182 varieties (out of 225) tested in all trials

Controls: Apache, Orvantis, Caphorn, and Soissons (07/08) or Premio (08/09)



Aerial view of EM08




Location

Genetic progress estimation



Additive effects

$$P_{ijk} = \mu + N_k + E_j + E_j \times N_k + \text{Gi} + \text{Gi} \times E_j + \text{Gi} \times N_k + \varepsilon_{ijk}$$



G = Quality + Precocity + Height + Year of Release


Quality classes	N°	YR means	GPC
High quality	104	2003 a	9.95 b
Good quality	50	2003 a	9.81 b
Biscuit quality	7	2001 ab	9.79 b
Other uses	7	2001 ab	9.20 c
Very high quality	12	1999 b	12.11 a



Genetic progress estimation

Interaction effect

$$P_{ijk} = \mu + N_k + E_j + E_j \times N_k + G_i + G_i \times E_j + G_i \times N_k + \varepsilon_{ijk}$$





$$\text{GxN} = \text{Quality} + \text{Precocity} + \text{Heigth} + \text{Year of Release}$$

Genetic progress estimation


Variance components

Trait	h^2	G	$G \times E$	$G \times N$
NupE	0.00	10% ns.	77% ***	13% ns.
NutE	0.79	63% ***	30% ***	7% ***
NUE	0.80	69% ***	26% ***	5% *



Genetic progress estimation

Interaction effects



Significant difference between
HN and LN



Gain in yield stability



Based on heritabilities and genetic correlations:
Direct selection at LN is still the more efficient
Indirect Selection Efficiency ISE=0.78

.010

How to maintain / accelerate this improvement?

- ✓ Increase genetic diversity
 - 2 new association panels (4600 INRA collection)
 - 10 advanced backcross populations



- ✓ Identify linked molecular markers

- ✓ Develop phenotyping tools



Association genetics parameters

214 genotyped varieties

34 382 polymorphic SNP (Illumina 90K + Biogemma)

Mean Linkage Disequilibrium (LD) decay = 1.1 +/- 1.3 cM


A poorly structured panel



Two first axes of the PCA of varieties by 3 461 non-redundant SNP

Example of an associated region


$$\text{NUE} = \text{ENV} + \text{Quality} + \text{Precocity} + \text{SNP} + \text{SNP} \times \text{N} + \text{K}$$



The most significant LOD score
of interaction with available N



Effect on remobilization (NHI
and %N in straw)



Region already identified in 2
independent yield GWAS
(# GW2)


.013

Epistasis and QTL6A


$$\text{NUE} = \text{ENV} + \text{Quality} + \text{Precocity} + \text{SNPi} \times \text{SNPj} + K$$

The effect of this region is significantly conditionned by others

20% of NUE genetic values can be explained by the interaction of QTL6A / Protein PSII




Interaction network based on significant epistatic interactions for NUE between SNP having a significant effect on NUE



Boxplot of NUE genetic values as a function of alleles at QTL6A/PSII

A possible candidate gene




.015

Looking for natural variants

Uauy et al. 2006 *TtNAM-A1* = IWGSC 6A seq



Genomic sequence of *TaNAM-A1*





Genotyping results of *TaNAM-A1*

GPC-A1 mapped in QTL6A confidence interval

→ polymorphism site

→ binding DNA aa



Theoretical 3D structure of natural variant of *TaNAM-A1* in European elite bread wheat (Chimera)

How to maintain / accelerate this improvement?

✓ Increase genetic diversity

- 2 new association panels (4600 INRA collection)
- 10 advanced backcross populations



✓ Identify linked molecular markers


✓ Develop phenotyping tools

Development of high-throughput techniques

- ✓ NUE is difficult to phenotype
 - time-consuming (sampling, milling ...)
 - not repeatable (biomass, N content)
- ✓ May explain that we detected no additive genetic diversity in the elite panel
- ✓ The key trait is total plant N
 - biomass and N content are difficult to estimate
 - probably possible to approximate them estimating Leaf Area (LAI) and Chlorophyll Content (CC)


Based on indices using reflectance values

LAI: MCARI2 - Modified Chlorophyll
Absorption Ratio Index 2




Haboudane et al., 2004

Chlorophyll: MTCI - MERIS
Terrestrial Chlorophyll Index



Dash and Curran, 2004

Developpement of a « PhenoMobile »



 **Effidence**

 **MECA 3D**  **UN PROJET DES SOLUTIONS**

 **ARVALIS**
Institut du végétal


 **INRA**
SCIENCE & IMPACT

F. Baret, B. de Solan, P. Burger


.020

Example of indices dynamics

Heading Flowering





MCARI2 : LAI related



MTCI : Chlorophylle related

Estimation of absorbed N



CONCLUSIONS

Estimation of genetic progress for NUE

- +0.33% / year
- significant G x N level interaction
- no significant genetic additive effect for N uptake

Association genetics in progress

- associated regions identified
- SNP x N level and SNP x SNP interactions detected

Phenotyping methods are currently developed

- LAI & Chlorophyll content dynamics

A second association panel was assembled (Breedwheat project)

- 220 elite lines, 100 common to the first panel
- genotyped (TaBW420k Affymetrix array, E. Paux et al)
- phenotyped
 - classical phenotyping (2012-2014)
 - high-throughput phenotyping (2012-2013)

.023

Acknowledgments

ANR ProtNBlé - S. Lafarge (Biogemma) et al.

ANR PhenoBlé - K. Beauchêne (Arvalis) et al.

IA Breedwheat - J. Le Gouis (INRA GDEC) et al.



BIOGEMMA :

Sébastien Praud (*Head « Genetic and Genomic of cereals » group*)

Fabien Cormier (PhD)

Mickael Throude

Pierre Dubreuil

Sébastien Faure

Stéphane Lafarge



Arvalis :

Katia Beauchêne

Benoit de Solan

Antoine Fournier



INRA:

Frédéric Baret

Philippe Burger

Alexis Comar

Emmanuel Heumez



Thank you for your attention

