



Identifying wild vs stocking components in fish recruitment despite the absence of identification data : an application to Atlantic salmon 0+ juveniles

G.J.R. Dauphin, Etienne Prévost

► To cite this version:

G.J.R. Dauphin, Etienne Prévost. Identifying wild vs stocking components in fish recruitment despite the absence of identification data : an application to Atlantic salmon 0+ juveniles. 4. International Statistical Ecology Conference (ISEC 2014), Jul 2014, Montpellier, France. 323 p. hal-02742526

HAL Id: hal-02742526

<https://hal.inrae.fr/hal-02742526>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Identifying wild vs stocking components in fish recruitment despite the absence of identification data: an application to Atlantic salmon 0+ juveniles.

G.J.R. Dauphin ^a and E. Prévost ^b

^a Department of Fisheries and Oceans
Northwest Atlantic Fisheries Center
80 East White Hills Road
PO Box 5667 St John's NL, Canada
Guillaume.Dauphin@dfo-mpo.gc.ca

^b INRA, UMR INRA-UPPA ECOBIOP
Aquapôle, Ibaron
64310 Saint-Pée sur Nivelle, France
eprevost@st-pee.inra.fr

Keywords: abundance; Atlantic salmon; hierarchical Bayesian model; juveniles; stocking; density dependence.

Abstract:

Over the last decades, exploited fish population have been declining significantly due to environmental changes, overexploitation and human activities. This has been reflected in the decline or collapse of numerous stocks. Alongside with fisheries regulations and habitat restoration, stock enhancement is a recurrent management tool used to inverse these trends. These enhancement programs are costly so measuring their impact is key although they are often poorly monitored.

During the course of an enhancement programme, stocking intensity may have been varied, intentionally or not, according to spatio-temporal units, wild production fluctuates in space and time too. The objective of this study is to propose an approach which takes advantage of these spatio-temporal variations of stocking vs wild recruitment for estimating their relative contribution from abundance data when no identification of the wild and stocked fish are available. We use Bayesian hierarchical modelling as a template for taking into account the spatio-temporal structure, while explicitly acknowledging that the abundance data issue from a mixture of fish of wild and hatchery origins. We illustrate our approach by means of a case study on 0+ juvenile of Atlantic salmon (*Salmo salar*) densities in the Allier catchment, France.

Not unlike similar stock enhancement projects in salmonids, the Allier stocking program lacks identification data for monitoring the fish that are released in the catchment separately from their wild counterparts. However, abundance data are available at the 0+ juvenile stage in early fall. A Bayesian hierarchical model was developed for estimating 0+ juvenile densities at two spatial scales over the 37 years of the study period, while assessing the relative contribution of wild reproduction, stocked eggs and stocked 0+ juvenile to the total 0+ juvenile production was evaluated.