



**HAL**  
open science

## The xyloglucans : are they new elicitors of *Arabidopsis thaliana* immunity ?

Justine Claverie, Daphnée Brulé, Xavier Daire, Benoît Poinssot

### ► To cite this version:

Justine Claverie, Daphnée Brulé, Xavier Daire, Benoît Poinssot. The xyloglucans : are they new elicitors of *Arabidopsis thaliana* immunity ?. 5. Journée des Doctorants de l'UMR 1347 Agroécologie, Institut National de la Recherche Agronomique (INRA). FRA., Mar 2016, Dijon, France. 27 p. hal-02742597

**HAL Id: hal-02742597**

**<https://hal.inrae.fr/hal-02742597v1>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# Journée des Doctorants

Lundi 14 Mars (9 h -14h)

Amphi Ampère – Bât. Gabriel

Au programme : 8 posters, 11 présentations orales dont 1 invité surprise + 1 buffet

Contacts : [carole.pfister@dijon.inra.fr](mailto:carole.pfister@dijon.inra.fr); [jeremie.zerbib@dijon.inra.fr](mailto:jeremie.zerbib@dijon.inra.fr)

## **The xyloglucans : are they new elicitors of *Arabidopsis thaliana* immunity ?**

**Claverie Justine**, Brulé Daphnée, Daire Xavier and Poinssot Benoit.

UMR Agroécologie 1347, INRA/Université de Bourgogne/Agrosup Dijon/CNRS, 17 rue Sully, BP 86510, 21000 Dijon, France

Damaged-Associated Molecular Patterns (DAMPs) are endogenous molecules released from the plant cell wall after wounding by pathogens. DAMPs are recognized by Pattern-Recognition Receptors (PRRs) that play a key role in plant immunity by mediating defense responses. The plant cell wall-derived oligogalacturonides (OG) are well characterized DAMPs that elicit plant immune responses such as MAPK activation, [Ca<sup>2+</sup>]<sub>cyt</sub> variations, H<sub>2</sub>O<sub>2</sub> production, defense-related gene expression and enhanced resistance against *Botrytis cinerea*. Our study focused on a new polysaccharide component of the plant cell wall called xyloglucans (Xh) and compared the immune events triggered by OG and Xh in *Arabidopsis thaliana*. Our results indicated that Xh can be considered as new elicitors as they induced MAPK activation, the expression of defense-related genes, callose deposition and triggered immunity against *Botrytis cinerea*. By using a genetic approach, our data indicated that the Xh-triggered immunity against *B. cinerea* requires the phytoalexin and jasmonic acid-dependent pathways.

Key words: Xyloglucans, Oligogalacturonides, DAMPs, Immunity, Signaling, *A. thaliana*.