



HAL
open science

Towards genome-wide association studies under abiotic stress in *Vitis vinifera*

Agnes Doligez, Loic L. Le Cunff, Timothée Flutre, Amandine Launay, Stephane S. Nicolas, Thierry Lacombe, Charles Romieu, Agota Fodor, Dalel Ahmed, Gilles Berger, et al.

► **To cite this version:**

Agnes Doligez, Loic L. Le Cunff, Timothée Flutre, Amandine Launay, Stephane S. Nicolas, et al.. Towards genome-wide association studies under abiotic stress in *Vitis vinifera*. ClimWine 2016 (Sustainable grape and wine production in the context of climate change), Apr 2016, Bordeaux, France. 152 p., 2016, Sustainable grape and wine production in the context of climate change. hal-02742772

HAL Id: hal-02742772

<https://hal.inrae.fr/hal-02742772v1>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Towards genome-wide association studies under abiotic stress in *Vitis vinifera*

A. Doligez¹, L. Le Cunff², T. Flutre¹, A. Launay¹, S. Nicolas³, T. Lacombe¹, C. Romieu¹,
A. Fodor², D. Ahmed¹, G. Berger¹, Y. Bertrand¹, M. Roques², P. This¹, J.-P. Péros¹

¹INRA, UMR AGAP, Montpellier, France

²IFV, UMT GenoVigne, Montpellier, France

³INRA, UMR GV, Gif-sur-Yvette, France

Genome-wide association studies have proven their efficiency to decipher the genetic determinism of traits of agronomic interest in several plant species. In order to perform such studies in grapevine, we first designed an association panel of 279 *V. vinifera* cultivars from the French National Grapevine Germplasm Collection (Domaine de Vassal, INRA, France). It is composed of three subgroups of 93 cultivars representing the three main genetic pools, which differ in use and geographical origin: wine West, wine East, table East. This panel exhibited a good genetic and phenotypic representativeness of the whole collection. Linkage disequilibrium extent was quite small, 43 Kb for a corrected r^2 predicted value of 0.2, emphasizing the need for several dozen thousands markers to achieve powerful genome wide association studies. The second step consisted in dense genotyping of this panel using a 18K SNP Infinium chip and genotyping by sequencing (GBS). While the first method provided useful genotypes at 12K SNPs, about 100K SNPs are expected using GBS with *ApeKI*. Data processing is in progress and this resource will be made available to the scientific community. The third step consists in phenotyping this panel for traits of interest. Vines were planted in a randomized block design at Le Chapitre Research Vineyard (Montpellier SupAgro, France). Traits related to yield (cluster and berry weights) and phenology (maturity date) were measured in three years within the DLVitis program. Then within the Innovine program, grass was used as a cover crop between rows in the whole trial and part of the design was irrigated to obtain two levels of water availability. In addition to yield and phenology traits, berry composition (sugars, acids, anthocyanins, tannins and aromas), pruning weight and adaptation to drought ($\delta C13$) have been measured during two growing seasons under both water availability conditions. This association panel, together with the accumulated data, represent highly valuable resources for genetic studies in grapevine. We are calling for an international collaborative effort to use such a panel in field and greenhouse trials to address questions on adaptation to changing environments.

Acknowledgments: INRA, DLVitis (ANR-08-GENM-02), UMT Geno-Vigne, Innovine (EU FP7 n° 31177).