



**HAL**  
open science

## **Interfacial behavior of milk polar lipids and their influence on gastric lipase adsorption: a natural effective delivery system**

Claire Bourlieu, Wafa Mahdoueni, Gilles Paboeuf, Samira de Oliveira, Stephane Pezenec, Jean-François Cavalier, Said Bouhallab, Didier Dupont, Pierre P. Villeneuve, Frédéric Carrière, et al.

### ► To cite this version:

Claire Bourlieu, Wafa Mahdoueni, Gilles Paboeuf, Samira de Oliveira, Stephane Pezenec, et al.. Interfacial behavior of milk polar lipids and their influence on gastric lipase adsorption: a natural effective delivery system. 107. AOCs Annual Meeting & Expo, May 2016, Salt Lake City, United States. hal-02743140

**HAL Id: hal-02743140**

**<https://hal.inrae.fr/hal-02743140>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# **Interfacial behavior of milk polar lipids and their influence on gastric lipase adsorption: a natural effective delivery system**

*C. Bourlieu<sup>1,2,5\*</sup>, W. Mahdoueni<sup>1,2</sup>, G. Paboeuf<sup>3</sup>, S. de Oliveira<sup>1,2</sup>, S. Pezennec<sup>1,2</sup>, J.-F. Cavalier<sup>4</sup>, S. Bouhallab<sup>1,2</sup>, D. Dupont<sup>1,2</sup>, P. Villeneuve<sup>5</sup>, F. Carrière<sup>4</sup>, V. Vié<sup>3</sup>*

*<sup>1</sup>INRA, UMR 1253 STLO, France, <sup>2</sup>Agrocampus Ouest UMR 1253 STLO, France; <sup>3</sup>IPR Institute of Physics, Rennes University 1, France, <sup>4</sup>CNRS, Aix-Marseille Université, UMR 7282 EIPL, France, <sup>5</sup>CIRAD, UMR IATE, France.*

The polar lipids in human milk fat globule membranes condition the accessibility and enzymatic digestibility of milk lipids. Their substitution by bovine polar lipids to produce infant formulas biomimetic of human milk has been suggested. However a comparison of the interfacial behavior of bovine and human polar lipids and of their interaction with the first enzyme involved in fat digestion, gastric lipase, is lacking.

Such comparison is here undertaken using complementary biophysical tools: tensiometry, ellipsometry, Brewster angle and atomic force microscopy, in the presence or in the absence of gastric lipase. Polar lipids extracts were obtained from a pool of human milk (n=5) or bovine butterserum and analyzed using GC and HPLC.

Human milk polar lipids (HMPL) presented a higher compressibility than bovine milk polar lipids (BMPL). Despite the presence of liquid condensed domains in both extracts, their morphological aspect and growth differed in relation with the lipid composition. Lipid phase separation impacted on gastric lipase adsorption in both extracts with an exclusive adsorption onto the liquid expanded phase. Despite differences in their physico-chemical properties, both polar lipid extracts share close interfacial reactivity in gastric conditions. This biophysical characterization will be broadened to other milk polar lipids interesting for human milk substitution.